

Searching for Diamonds

Jerrold R. Griggs
University of South Carolina

December 2, 2011

Abstract

Given a finite poset P , we consider the largest size $\text{La}(n, P)$ of a family of subsets of $[n] := \{1, \dots, n\}$ that contains no (weak) subposet P . Letting P_k denote the k -element chain (path poset), Sperner's Theorem (1928) gives that $\text{La}(n, P_2) = \binom{n}{\lfloor n/2 \rfloor}$, and Erdős (1945) showed more generally that $\text{La}(n, P_k)$ is the sum of the k middle binomial coefficients in n . Gyula Katona and his collaborators obtained many significant results for other posets P ; these results lead to the conjecture that $\pi(P) := \lim_{n \rightarrow \infty} \text{La}(n, P) / \binom{n}{\lfloor n/2 \rfloor}$ exists for general posets P , and in fact it is an integer.

For $k \geq 2$ let D_k denote the k -diamond poset $\{A < B_1, \dots, B_k < C\}$. By bounding the average number of times a random full chain meets a P -free family \mathcal{F} , called the Lubell function of \mathcal{F} , we prove that $\pi(D_2) < 2.273$, if it exists. This is a stubborn open problem, since we expect $\pi(D_2) = 2$. It is then surprising that, with appropriate partitions of the set of full chains, we can explicitly determine $\pi(D_k)$ for infinitely many values of k , and, moreover, describe the extremal D_k -free families. For these fortunate values of k , and for a growing collection of other posets P , we have that $\text{La}(n, P)$ is a sum of middle binomial coefficients in n , while for other values of k and for most P , it seems that $\text{La}(n, P)$ is far more complicated.

This is joint work with Wei-Tian Li and Linyuan Lu.