

Charge Transfer Enhances Two-Photon Absorption in Transition Metal Porphyrins

Jonathan L. Humphrey and Darius Kuciauskas*

Department of Chemistry, Virginia Commonwealth University, 1001 West Main Street,
Richmond, Virginia 23284-2006

Received January 9, 2006; E-mail: dkuciauskas@gmail.com

Porphyrin-based materials are promising for applications that require strong two-photon absorption.¹ Such applications include fluorescence microscopy and photodynamic therapy with two-photon excitation, 3D microfabrication, and sensor protection.¹ Enhancement of the optical properties that determine two-photon absorption efficiency (molecular second hyperpolarizability, γ , and two-photon absorption cross-section, δ) have been achieved through modifications to the π network of the porphyrin macrocycle.¹ In this report, we consider the effect of transition metal ions on the porphyrin γ and δ values.² We show that metalloporphyrins with strong charge transfer (CT) transitions in their one-photon absorption spectra have δ values that are an order of magnitude larger than those of the free base porphyrin. We attribute this effect to CT transitions between the metal ion's d orbitals and the π system of the porphyrin.

Electropolymerized thin films of Fe³⁺,³ Mn³⁺, and Co²⁺ 5,10,15,20-tetrakis-(4-hydroxytetraphenyl)porphyrins (TPP) were studied. Because these metalloporphyrins have low emission yields, degenerate four wave mixing (DFMW) experiments with 100 fs laser pulses were used to determine γ and δ . Details of the film preparation and optical studies are given in the Supporting Information.

The absorption spectra of Fe³⁺, Mn³⁺, and Co²⁺ TPP are shown in Figure 1A. The widely used four orbital model (a_{1u}, a_{2u} HOMO and e_g LUMO, Figure 1B) qualitatively explains the origin of porphyrin B and Q bands.⁴ For Fe³⁺ and Mn³⁺ TPP, mixing of the metal d and macrocycle π orbitals leads to additional peaks in the absorption spectrum.^{5,6} For Fe³⁺ TPP, these CT transitions involve the π (a_{1u}, a_{2u}) orbitals of the macrocycle and the metal ion's d_{π} orbitals ($\pi \rightarrow d_{\pi}$).⁶ For Mn³⁺ TPP, the CT bands include the π (b_{2u}) $\rightarrow d_{\pi}$ ($\lambda_{\max} = 432$ nm) and π (a_{1u}, a_{2u}) $\rightarrow d_{z^2}$ ($\lambda_{\max} = 638$ nm) transitions.⁶ These CT transitions are allowed as they occur between electronic states of opposite symmetry ($g \rightarrow u$). The absence of similar CT bands in the Co²⁺ TPP absorption spectrum suggests that the mixing of the Co²⁺ ion's d orbitals with the macrocycle's π system is less efficient. Thus, comparison of Co²⁺ TPP and Mn³⁺/Fe³⁺ TPP optical properties allows the role of metal ion/macrocycle interactions on two-photon effects to be investigated.

The metalloporphyrin second hyperpolarizability, γ , spectra are shown in Figure 2. (DFWM data are provided in Supporting Information. The ~ 100 fs response time of each material indicates a purely electronic process.) Peaks are observed in the near-IR region of the γ spectra at ~ 1150 nm for Fe³⁺ TPP and ~ 1250 nm for Mn³⁺ TPP. The Co²⁺ TPP γ spectrum does not have this feature. The molecular orbital model shown in Figure 1B suggests that the peaks in the γ spectra could be related to two-photon transitions between metal d and macrocycle π^* orbitals. As both orbitals have g symmetry, two-photon transitions are allowed. In contrast, one-photon transitions between these orbitals are forbidden and are not seen in the absorption spectra in Figure 1A.




Figure 1. (A) One-photon absorption spectrum of Mn³⁺ (—), Fe³⁺ (—), and Co²⁺ (—) TPP. (B) Molecular orbital diagram for Mn³⁺ and Fe³⁺ TPP. Metal orbitals are shown in red. Adapted from refs 11–13




Figure 2. Second hyperpolarizability, γ , of (A) Mn³⁺, (B) Fe³⁺, and (C) Co²⁺ TPP determined by DFWM studies on thin films. The solid lines are fitting results according to eq 1. The two-photon component ($\text{Im } \gamma$, dashed red line) is also shown.

A sum-over-states model was used to estimate the two-photon state energies and dipole moments. A three-state model (ground, one-, and two-photon excited states) was previously applied to porphyrin materials:⁷

$$\gamma \propto -\frac{M_{01}^4}{(\omega_1 - \omega - i\Gamma)^3} + \frac{M_{01}^2 M_{12}^2}{(\omega_1 - \omega - i\Gamma)^2 (\omega_2 - 2\omega - i\Gamma)} \quad (1)$$

where M_{01} and M_{12} are the transition dipole moments for the $0 \rightarrow 1$ and $1 \rightarrow 2$ transitions, ω is the experimental frequency, ω_1 and ω_2 are frequencies derived from the excited-state energies E_1 and E_2 ($\omega_1 = E_1/h$ and $\omega_2 = E_2/h$; h is the reduced Planck's constant). Parameter Γ accounts for the linewidths. Analysis of the γ spectra yields the M_{12} , E_2 , and Γ values (Table 1; E_1 and M_{01} were obtained from absorption spectra in Figure 1A). As suggested by the γ spectra in Figure 2, the Fe³⁺ TPP two-photon state has a higher energy than that of Mn³⁺ TPP. Both porphyrins have M_{12} values of 4.2 D.

Table 1. Linear and Nonlinear Spectroscopic Properties of the Metalloporphyrin Films

	$M_{01}(\text{D})^a$	$M_{12}(\text{D})^b$	$E_1(\text{cm}^{-1})^a$	$E_2(\text{cm}^{-1})^b$	$\Gamma(\text{cm}^{-1})^a$
Mn^{3+}	3.85	4.2	15686	16050	1150
Fe^{3+}	1.3	4.2	14300	17400	900

^a The errors are $\pm 3.5\%$ in these values. ^b $\pm 10\%$ error.




Figure 3. One- (—) and two-photon (---) spectra of Mn^{3+} TPP (black) and Fe^{3+} TPP (red).

To determine the effect of the metalloporphyrin two-photon states on the two-photon absorption cross-section, δ , we consider the imaginary component of hyperpolarizability, $\text{Im } \gamma$. The cross-section δ can be obtained from:⁸

$$\delta = \frac{6\pi^2 \hbar}{\epsilon_0 n^2 \lambda^2} f^4 \text{Im } \gamma \quad (2)$$

where ϵ_0 is the electric constant, n is the index of refraction, and λ is the wavelength. Application of eq 2 to the $\text{Im } \gamma$ data yields $\delta_{\text{max}} = 82 \text{ GM}$ for Fe^{3+} TPP and $\delta_{\text{max}} = 164 \text{ GM}$ for Mn^{3+} TPP (Figure 3; $1 \text{ GM} = 1 \times 10^{-50} \text{ cm}^4 \text{ s}^{-1}$). The difference in δ_{max} values can be attributed to the increase in the transition dipole moment, M_{01} , of Mn^{3+} TPP over that of Fe^{3+} TPP. For metal-free TPP, $\delta < 10 \text{ GM}$ in the spectral range of our experiments.⁹ Therefore, Mn^{3+} and Fe^{3+} ions enhance two-photon absorption intensity by at least an order of magnitude.

A comparison of the one- and two-photon spectra (Figure 3, the energy scale in the upper axis reflects a two-photon process) offers further support for the assignment of the two-photon absorption bands (dashed lines) to intramolecular charge transfer. The two-photon transitions do not originate in the π network of the macrocycle, as they are not allowed by the selection rules. If the selection rules for the two-photon processes are relaxed due to symmetry breaking from distortion of the macrocycle in the solid state, two-photon absorption would be stronger into the vibronic Q(0,1) band than into the purely electronic Q(0,0) band.⁹ Our results show the opposite trend: two-photon absorption is stronger into the region of the Q(0,0) band than into that of the Q(0,1) band. Additionally, the porphyrin macrocycle's two-photon states were predicted to have much higher energies than those observed in our experiments.¹⁰

Shifts in the energies of the two-photon states, E_2 , suggest which molecular orbitals give rise to the two-photon CT transitions (according to the four orbital model, Figure 1B). For Fe^{3+} and Mn^{3+} TPP, the metal d_{xy} , d_{π} , and d_z orbitals are in the HOMO/LUMO gap of the macrocycle's π orbitals. The Fe^{3+} ion's d_{xy} orbital is very close in energy to the a_{1u}, a_{2u} orbitals,¹¹ and the Fe^{3+} TPP two-photon state energy determined in our experiments is very close in energy to the Q(0,0) state. This suggests involvement of the d_{xy} orbital in the two-photon transition, $d_{xy} \rightarrow \pi^*$. For the Mn^{3+} ion,

the d_{xy} orbital is at a higher energy than the macrocycle's a_{1u}, a_{2u} orbitals.¹¹ If the two-photon transition originates from the d_{xy} orbital, the Mn^{3+} TPP state will be at a lower energy than that observed for Fe^{3+} TPP. The experimental results agree with this prediction (Table 1 and Figure 3). Definite assignment of the molecular orbitals involved in the two-photon transition will require quantum chemical computations for molecular excited states. Such studies are difficult for transition metal complexes with an unfilled d electron shell.¹²

While the Co^{2+} d_{xy} orbital is at an energy similar to those of the Mn^{3+} and Fe^{3+} ions,¹³ we did not observe CT transitions in Co^{2+} TPP γ spectra. The much weaker transition dipole moment, M_{01} , of this metalloporphyrin could account for this result. While the CT absorption bands have $\epsilon = 2300 \text{ M}^{-1} \text{ cm}^{-1}$ for Fe^{3+} TPP and $\epsilon = 10000 \text{ M}^{-1} \text{ cm}^{-1}$ for Mn^{3+} TPP, the corresponding extinction coefficient of Co^{2+} TPP was only $\epsilon \approx 75 \text{ M}^{-1} \text{ cm}^{-1}$.¹⁴ The small ϵ value implies that the M_{01} of Co^{2+} TPP is too small to promote amplification of the two-photon processes. The two-photon absorption enhancement in linear conjugated oligomers is also dependent on M_{01} .¹⁵

In summary, Fe^{3+} and Mn^{3+} ions enhance the second hyperpolarizability, γ , and two-photon absorption cross-section, δ , of transition metal porphyrins by at least an order of magnitude. The enhancement is dependent on the one-photon transition dipole moment. A two-photon $d_{xy} \rightarrow \pi^*$ charge-transfer transition was observed for Fe^{3+} and Mn^{3+} tetraphenyl porphyrins for the first time, and suggest a new approach to enhance optical properties of porphyrin materials.

Acknowledgment. Support from VCU is gratefully acknowledged.

Supporting Information Available: Metalloporphyrin film characterization, DFWM kinetics and spectra, data analysis used to obtain γ spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- Ogawa, K.; Ohashi, A.; Kobuke, Y.; Kamada, K.; Ohta, K. *J. Am. Chem. Soc.* **2003**, *125*, 13356; Drobizhev, M.; Stepanenko, Y.; Dzenis, Y.; Karotki, A.; Rebane, A.; Taylor, P. N.; Anderson, H. L. *J. Am. Chem. Soc.* **2004**, *126*, 15352; Screen, T. E. O.; Thorne, J. R. G.; Denning, R. G.; Bucknall, D. G.; Anderson, H. L. *J. Am. Chem. Soc.* **2002**, *124*, 9712.
- Screen, T. E. O.; Thorne, J. R. G.; Denning, R. G.; Bucknall, D. G.; Anderson, H. L. *J. Mater. Chem.* **2003**, *13*, 2796; Kandasamy, K.; Shetty, S. J.; Puntambekar, P. N.; Srivastava, T. S.; Kundu, T.; Singh, B. P. *J. Porphyrins Phthalocyanines* **1999**, *3*, 81.
- Humphrey, J.; Kuciauskas, D. *J. Phys. Chem. B* **2004**, *108*, 12016.
- Gouterman, M. *J. Chem. Phys.* **1959**, *30*, 1139.
- Kobayashi, H.; Yanagawa, Y.; Osada, H.; Minami, S.; Shimizu, M. *Bull. Chem. Soc. Jpn.* **1973**, *46*, 1471; Ohya, T.; Sato, M. *Bull. Chem. Soc. Jpn.* **1996**, *69*, 3201.
- Boucher, L. J. *J. Am. Chem. Soc.* **1970**, *92*, 2725.
- Thorne, J. R. G.; Kuebler, S. M.; Denning, R. G.; Blake, I. M.; Taylor, P. N.; Anderson, H. L. *J. Chem. Phys.* **1999**, *248*, 181.
- Gubler, U.; Bosshard, C. In *Polymers in Photonics Applications I*; Lee, K.-S., Ed.; Springer: Berlin, 2002; p 123.
- Karotki, A.; Drobizhev, M.; Kruk, M.; Spangler, C.; Nickel, E.; Mamedashvili, N.; Rebane, A. *J. Opt. Soc. Am. B* **2003**, *20*, 321; Kruk, M.; Karotki, A.; Drobizhev, M.; Kuzmitsky, V.; Gael, V.; Rebane, A. *J. Lumin.* **2003**, *105*, 45.
- Masthay, M. B.; Finsden, L. A.; Pierce, B. M.; Bocian, D. F.; Lindsey, J. S.; Birge, R. R. *J. Chem. Phys.* **1986**, *84*, 3901; Kuzmitsky, V. A. *J. Appl. Spectrosc.* **2001**, *68*, 758; Liu, X. J.; Feng, J. K.; Ren, A. M.; Zhou, X. *Chem. Phys. Lett.* **2003**, *373*, 197; Zhou, X.; Ren, A. M.; Feng, J. K.; Liu, X. J.; Zhang, Y. D. *ChemPhysChem* **2003**, *4*, 991.
- Gouterman, M.; Hanson, L. K.; Khalil, G. E.; Leenstra, W. R.; Buchler, J. W. *J. Chem. Phys.* **1975**, *62*, 2343.
- Liao, M.-S.; Watts, J. D.; Huang, M.-J. *J. Phys. Chem. A* **2005**, *109*, 11996.
- Zerner, M.; Gouterman, M. *Theor. Chim. Act.* **1966**, *4*, 44.
- Antipas, A.; Gouterman, M. *J. Am. Chem. Soc.* **1983**, *105*, 4896.
- Rumi, M.; Ehrlich, J. E.; Heikal, A. A.; Perry, J. W.; Barlow, S.; Hu, Z. Y.; McCord-Maughon, D.; Parker, T. C.; Rockel, H.; Thayumanavan, S.; Marder, S. R.; Beljonne, D.; Bredas, J. L. *J. Am. Chem. Soc.* **2000**, *122*, 9500.

JA0588353