

Pebbling in powers of paths

Liliana Alc3n *

Glenn Hurlbert †

December 2, 2022

Dedicated to the memory of Ron Graham.

Abstract

The t -fold pebbling number, $\pi_t(G)$, of a graph G is defined to be the minimum number m so that, from any given configuration of m pebbles on the vertices of G , it is possible to place at least t pebbles on any specified vertex via pebbling moves. It has been conjectured that the pebbling numbers of pyramid-free chordal graphs can be calculated in polynomial time.

The k^{th} power $G^{(k)}$ of the graph G is obtained from G by adding an edge between any two vertices of distance at most k from each other. The k^{th} power of the path P_n on n vertices is an important class of pyramid-free chordal graphs, and is a stepping stone to the more general class of k -paths and the still more general class of interval graphs. Pachter, Snevily, and Voxman (1995) calculated $\pi(P_n^{(2)})$, Kim (2004) calculated $\pi(P_n^{(3)})$, and Kim and Kim (2010) calculated $\pi(P_n^{(4)})$. In this paper we calculate $\pi_t(P_n^{(k)})$ for all n , k , and t .

For a function $D : V(G) \rightarrow \mathbb{N}$, the D -pebbling number, $\pi(G, D)$, of a graph G is defined to be the minimum number m so that, from any given configuration of m pebbles on the vertices of G , it is possible to place at least $D(v)$ pebbles on each vertex v via pebbling moves. It has been conjectured that $\pi(G, D) \leq \pi_{|D|}(G)$ for all G and D . We make the stronger conjecture that every G and D satisfies $\pi(G, D) \leq \pi_{|D|}(G) - (s(D) - 1)$, where $s(D)$ counts the number of vertices v with $D(v) > 0$. We prove that trees and $P_n^{(k)}$, for all n and k , satisfy the stronger conjecture.

The pebbling exponent $e_\pi(G)$ of a graph G was defined by Pachter, et al., to be the minimum k for which $\pi(G^{(k)}) = n(G^{(k)})$. Of course, $e_\pi(G) \leq \text{diam}(G)$, and Czygrinow, Hurlbert, Kierstead, and Trotter (2002) proved that almost all graphs G have $e_\pi(G) = 1$. Lourdasamy and Mathivanan (2015) proved several results on $\pi_t(C_n^2)$, and Hurlbert (2017) proved an asymptotically tight formula for $e_\pi(C_n)$. Our formula for $\pi_t(P_n^{(k)})$ allows us to compute $e_\pi(P_n)$ within additively narrow bounds.

*CeMaLP, UNLP, CONICET, Argentina

†Department of Mathematics and Applied Mathematics, Virginia Commonwealth University, USA

2000 AMS Subject Classification: 05C40, 05C75, 05C87 and 05C99.

Keywords: graph pebbling, pebbling number, pebbling exponent, target conjecture.

1 Introduction

Graph pebbling has an interesting history, with many challenging open problems. Calculating pebbling numbers of graphs is a well known computationally difficult problem (in Π_2^P -complete [19]). See [11, 13] for more background. It has been asked (e.g. [12]) for what families of graphs G the pebbling number $\pi(G)$ (defined below) can be calculated in polynomial time. One possible family posited in [1] is that of chordal graphs, most likely with some restriction, such as bounded diameter or treewidth, for example. This paper follows a sequence ([1, 2, 3, 4]) intended to provide at least a partial answer to this line of inquiry, which has led us to make the following conjecture. We define the *pyramid* to be the triangulated 6-cycle $abcdef$ with interior triangle bdf , and say that a graph is *H-free* if it does not contain H as an induced subgraph.

Conjecture 1. *If G is a pyramid-free chordal graph then $\pi(G)$ can be calculated in polynomial time.*

A *configuration* C of pebbles on the vertices of a connected graph G is a function $C : V(G) \rightarrow \mathbb{N}$ (the nonnegative integers), so that $C(v)$ counts the number of pebbles placed on the vertex v . A vertex v is *empty* if $C(v) = 0$ and *big* if $C(v) \geq 2$. We write $|C|$ for the *size* $\sum_v C(v)$ of C ; i.e. the number of pebbles in the configuration. A *pebbling step* from a big vertex u to one of its neighbors v (denoted $u \rightarrow v$) reduces $C(u)$ by two and increases $C(v)$ by one. Given a specified *target* vertex r we say that C is *t-fold r-solvable* if some sequence of pebbling steps places t pebbles on r . We are concerned with determining $\pi_t(G, r)$, the minimum positive integer m such that every configuration of size m on the vertices of G is *t-fold r-solvable*. The *t-pebbling number* of G is defined to be $\pi_t(G) = \max_{r \in V(G)} \pi_t(G, r)$. We adopt the natural interpretation that $\pi_0(G) = 0$, and avoid writing t when $t = 1$.

The k^{th} power $G^{(k)}$ of the graph G is obtained from G by adding an edge between any two vertices of distance at most k from each other. The *pebbling exponent* $e_\pi(G)$ of a graph G was defined in [21] to be the minimum k for which $\pi(G^{(k)}) = n(G^{(k)})$. For example, *Class 0* graphs (graphs G with $\pi(G) = n(G)$) have pebbling exponent $e_\pi(G) = 1$. In a very strong probabilistic sense (see [8]) almost all graphs G have $e_\pi(G) = 1$. Of course, $e_\pi(G) \leq \text{diam}(G)$, and the authors of [21] ask specifically about the cycle C_n on n vertices. In [14] it was shown that $n/(2 \lg n) \leq e_\pi(C_n) \leq n/[2(\lg n - \lg \lg n)]$, which determines its exact value for $n \leq 9$, bounds it within a factor of two always, and a factor of one asymptotically. Here we write \lg for the base 2 logarithm. Lourdasamy and Mathivanan [18] proved several results on $\pi_t(C_n^2)$.

Denote the path on n vertices by P_n . Pachter, et al. [21] proved that $\pi(P_n^{(2)}) = 2^{\lceil \frac{n-1}{2} \rceil} + ((n-2) \bmod 2)$ for $n \geq 2$. Kim [16] proved that $\pi(P_n^{(3)}) = 2^{\lceil \frac{n-1}{3} \rceil} + ((n-2) \bmod 3)$ for $n \geq 8$ (and equals n for $n \leq 7$). Kim and Kim [17] proved that $\pi(P_n^{(4)}) = 2^{\lceil \frac{n-1}{4} \rceil} + ((n-2) \bmod 4)$ for $n \geq 14$ (and equals n for $n \leq 13$). We generalize these results in Theorem 6, below, calculating $\pi_t(P_n^{(k)})$ for all k and all t . This allows us to compute $e_\pi(P_n)$ very tightly in Corollary 8.

We generalize the traditional pebbling model as follows. A *pebbling function* F is any function $F : V \rightarrow \mathbb{N}$; its *size* is $|F| = \sum_{v \in V} F(v)$. For a pebbling function F , define \dot{F} to be the multiset $\{v^{F(v)}\}_{v \in V}$ (the exponent $F(v)$ is the *multiplicity* of v). Configurations C (of pebbles) and distributions D (of targets) are both pebbling functions. However, we think of \dot{C} as a multiset of pebbles, labeled by their vertex locations, while we think of \dot{D} as a multiset of target vertices. Furthermore, we think of $v_{i,j}$ as the label of j^{th} pebble (or target) sitting on vertex v_i . For $m \in \mathbb{N}$ and $x \in V$, define the function mx by $mx(v) = m$ if $v = x$ and 0 otherwise. Thus, the symbol x can represent a vertex or a pebbling function, depending on its context. In particular, if $F = mx$ then $\dot{F} = \{x^m\}$.

For a configuration C and distribution D , we say that C is D -solvable (or that there is a (C, D) -solution, or that G has a (C, D) -solution) if some sequence of pebbling steps places at least $D(v)$ pebbles on each vertex v . The D -pebbling number, $\pi(G, D)$, of a graph G is defined to be the minimum number m such that G is (C, D) -solvable whenever $|C| \geq m$. Thus we can write $\pi_t(G, r)$ as $\pi(G, tr)$ in this generalized notation. (We note that the D -pebbling number was first introduced in [7] for the case $D = V(G)$, and was called the *cover pebbling number*.)

Just as the t -fold pebbling number can be used inductively to prove results about the pebbling number (e.g. [6, 3]), the D -pebbling number can be used inductively to prove results about the t -fold pebbling number. It is also thought that this might be a powerful tool in attacking Graham's Conjecture on the pebbling number of the cartesian product of graphs (see [6]). The following **Weak Target Conjecture** was conjectured in [9].

Conjecture 2. [9] *Every graph G satisfies $\pi(G, D) \leq \pi_{|D|}(G)$ for every target distribution D .*

The authors of [9] verified this conjecture for trees, cycles, complete graphs, and cubes, and the authors of [15] verified this conjecture for 2-paths and Kneser graphs $K(m, 2)$ with $m \geq 5$.

Define $\text{supp}(D)$ to be the set of vertices v with $D(v) > 0$, and denote $s(D) = |\text{supp}(D)|$. We make the following **Strong Target Conjecture**.

Conjecture 3. *Every graph G satisfies $\pi(G, D) \leq \pi_{|D|}(G) - (s(D) - 1)$ for every target distribution D .*

We prove that trees satisfy this stronger conjecture in Theorem 4. We prove in Theorem 5 that $P_n^{(k)}$ satisfies this stronger conjecture for all n and k .

Another reason to study pebbling on powers of paths is the following. Czygrinow, et al. [8], proved that, for each $d \geq 1$, there is a least positive integer $k(d)$ such that if G has diameter d and connectivity at least $k(d)$ then G is Class 0. They showed that $k(d) \leq 2^{2d+3}$ and $k(d) \in \Omega(2^d/d)$. We note that $P_n^{(k)}$ has connectivity equal to k and that Theorem 6 shows that $\pi(P_n^{(k)}) = n$ when $k \geq (2^d - 2)/(d - 1)$. Thus $P_n^{(k)}$

witnesses the tightness of the lower bound on $k(d)$; we believe that the upper bound on $k(d)$ is weak and should be improved. Furthermore, at the other extreme when t is large, a theorem of [10] states that every graph G satisfies $\lim_{t \rightarrow \infty} \pi_t(G)/t = 2^d$, where $d = \text{diam}(G)$. Our Theorem 6 formula for $t \geq k(d-1)/(2^d - 2)$ is more precise, that $\pi_t(P_n^{(k)}) = t2^d + ((n-2) \bmod k)$.

The final and, for our current purposes, most important motivation for investigating $\pi_t(P_n^{(k)})$ for $2 \leq k \leq \text{diam}(P_n^{(k)}) = \lfloor (n-2)/k \rfloor + 1$ is that $P_n^{(k)}$ is chordal. This case is a key stepping stone toward the graph classes k -paths and interval graphs. At each stage of the sequence of papers [1, 2, 3, 4] mentioned above we have discovered new hurdles that have required new techniques which have expanded our understanding of pebbling in chordal graphs, such as the technical lemmas found in Section 3. The critical pieces of the puzzle in this paper are the new chordal lemmas found in Section 3.3, as well as the careful interplay between the t -wide and t -long cases in the inductive proof of Section 5. In particular, Conjecture 3 plays a crucial role and may be the most important contribution of this work, as a powerful technique in future research.

We describe our results in the next section, introduce the important machinery in Section 3, and prove Theorems 4–6 in order in Sections 4–6.

2 Main Theorems

For positive integers n and k , the *path power* $P_n^{(k)}$ is the graph with vertex set $V = \{v_1, \dots, v_n\}$, and $v_i \sim v_j$ whenever $1 \leq |i - j| \leq k$. The diameter d of $P_n^{(k)}$ is completely determined by n and k ; in fact,

$$d = \lfloor (n-2)/k \rfloor + 1.$$

We let $b = (n-2) - k(d-1)$, then $0 \leq b < k$. For $n \geq 2$ and $t \geq 1$, define the following functions, where the above formula for d is assumed:

$$\begin{aligned} l_t(n, k) &= t2^d + n - 2 - k(d-1) = t2^d + b, \\ w_t(n) &= 2t + n - 2, \\ p_t(n, k) &= \max\{l_t(n, k), w_t(n)\}. \end{aligned}$$

We have $w_t(n) \geq l_t(n, k)$ if and only if $t(2^d - 2) \leq (d-1)k$; in other words

$$p_t(n, k) = \begin{cases} 2t + n - 2 & \text{if } t \leq t_0, \text{ and} \\ t2^d + b & \text{if } t \geq t_0, \end{cases}$$

where $t_0 = 1$ if $d = 1$, and $t_0 = t_0(k, d) = k(d-1)/(2^d - 2)$ otherwise. We say that $P_n^{(k)}$ is *t-wide* when $t \leq t_0$ and *t-long* when $t \geq t_0$. More finely, we say that $P_n^{(k)}$ is *barely t-long* when $t_0 < t = \lceil t_0 \rceil$, and *strictly*

t -long if $t > \lceil t_0 \rceil$. Thus, the formula names w_t and l_t match the t -wide and t -long terminology. We will occasionally make use of the fact that, for $d \geq 2$ and fixed k , $t_0(k, d)$ is a strictly decreasing function of d .

For $k \geq n-1$, $P_n^{(k)}$ is a complete graph, and it is t -wide for any $t \geq 1$. Therefore, $\pi_t(P_n^{(k)}, r) = \pi_t(P_n^{(k)}) = 2t + n - 2 = w_t(n) = p_t(n, k)$ for any vertex r . Notice that in this case all vertices are *simplicial* (i.e. their neighborhoods induce complete graphs). For $1 \leq k < n-1$, the vertices v_1 and v_n are the only two simplicial vertices of $P_n^{(k)}$; the non simplicial vertices will be called *interior* vertices. Note that the connectivity of $P_n^{(k)}$ equals k . Moreover, when $1 \leq (n-1)/2 \leq k$, $P_n^{(k)}$ contains a universal vertex (adjacent to every other vertex); then the present work extends that of [4], which computes the pebbling numbers of k -connected graphs with universal vertices. For practical purposes, when $n = 1$, we define $p_t(1, k) = t - 1$ for any k .

The next three theorems will be proved in Sections 4–6.

The following theorem verifies the Strong Target Conjecture for trees. The cost of a solution refers to the total number of pebbles that are lost when performing the steps of the solution, the formal definition is given in the next section.

Theorem 4. *Let T be a tree of diameter d and D be a target distribution of size t . Then $\pi(T, D) \leq \pi_t(T) - s(D) + 1$. Furthermore, if C is a configuration on T of size $|C| \geq \pi_t(T) - s(D) + 1$, then C solves any target $v \in \dot{D}$ with cost at most $2^{\text{ecc}_T(v)}$.*

The following theorem verifies the Strong Target Conjecture for powers of paths.

Theorem 5. *Let D be a t -multiset of target vertices of $P_n^{(k)}$. Then $\pi(P_n^{(k)}, D) \leq p_t(n, k) - s(D) + 1$.*

Analogous to Theorem 4, as part of the proof of Theorem 5, we show in the t -long case that if C is a configuration on $G = P_n^{(k)}$ of size $|C| \geq p_t(n, k) - s(D) + 1$ then C solves any target $v \in \dot{D}$ with cost at most $2^{\text{ecc}_G(v)}$.

Theorem 6. $\pi_t(P_n^{(k)}) = p_t(n, k)$.

As a result we obtain the following two corollaries.

Corollary 7. *The t -fold pebbling number $\pi_t(P_n^{(k)})$ can be calculated in polynomial time (constant time if k is known and linear time otherwise).*

The constant time follows from simply comparing $t(2^d - 2)$ with $k(d - 1)$, while the linear time follows from knowing that $k = \delta(P_n^{(k)})$ (which equals the degree of a simplicial vertex).

Corollary 8. *Define the functions $M(n) = \lfloor (n-2)/(\lceil \lg n \rceil - 2) \rfloor$ and $m(n) = \lfloor (n-2)/(\lceil \lg n \rceil - 2)^2 \rfloor$. Then the pebbling exponents of paths are $e_\pi(P_2) = 1$, $e_\pi(P_3) = e_\pi(P_4) = e_\pi(P_5) = 2$, $e_\pi(P_6) = e_\pi(P_7) = e_\pi(P_8) = 3$, and, for $n \geq 9$, $M(n) - m(n) \leq e_\pi(P_n) \leq M(n)$.*

We prove this in Section 7. We also note that the upper bound is tight at $n = 33, 65, 257$, and many other values, typically of the form $n = 2^i + j$ for a few small values of j . In fact, numerical evidence suggests that it may be tight at $n = 2^i + 1$ for most values of $i \geq 5$. Moreover, except for values of n in the range of something like $[2^i, 2^i + i^2)$, it appears that the tighter bounds $M(n) - m(n) + 2 \leq e_\pi(P_n) \leq M(n) - m(n) + 3$ may hold.

3 Technical Lemmas

3.1 General Lemmas

Given a configuration C of pebbles, a *potential move* is either a pair of pebbles sitting on the same vertex, or a single pebble sitting on a target vertex, which in either case is called *potential vertex*. When counting the number of potential moves of C in relation to a target D , we must be careful about counting too many singletons on a target vertex v : $\min\{C(v), D(v)\}$ of the pebbles there are singleton potentials, while the other $C(v) - D(v)$ pebbles must be counted in pairs (since they would need to move to solve other targets). To say that C has j potential moves means that the j pairs and singletons are pairwise disjoint. For example, the configuration C on 5 vertices (v_1, \dots, v_5) with values $(0, 1, 1, 2, 7)$ has 4 potential moves if the target D has values $(2, 0, 0, 0, 0)$, and 5 potential moves if D has values $(1, 0, 0, 0, 2)$. The *potential* of C , $\text{pot}(C)$, is the maximum j for which C has j potential moves. Because every solution that requires a pebbling move uses a potential move, the following fact is evident.

Fact 9. *Let r be an empty vertex in a configuration C with $\text{pot}(C) < t$. Then C is not t -fold r -solvable.*

Another useful tool is the following lemma.

Lemma 10. (Potential Lemma) *Let G be a graph on n vertices. If C is a configuration on G of size $n + y$ ($y \geq 0$) having z zeros, then $\text{pot}(C) \geq \lceil \frac{y+z}{2} \rceil$.*

A (*induced*) *slide* from a potential vertex v to a vertex r is a (induced) path between v and r with a pebble on each interior vertex. Two slides are disjoint if the corresponding sets of pebbles are disjoint.

Let C be a configuration of c pebbles p_1, p_2, \dots, p_c on a graph G . Let \mathcal{S} be a t -fold r -solution of C moving pebbles p_1, p_2, \dots, p_t into the target vertex r . This means that \mathcal{S} is a sequence $\sigma_1, \sigma_2, \dots, \sigma_h$ of pebbling steps after which a configuration with the pebbles p_1, p_2, \dots, p_t on r is obtained. Assume that σ_i moves the pebble p_{σ_i} from vertex v_{σ_i} to vertex v'_{σ_i} and discards the pebble p'_{σ_i} . We define the directed multigraph $G(\mathcal{S})$ to have the same vertex set as G , with a directed edge $(v_{\sigma_i}, v'_{\sigma_i})$ for each pebbling step σ_i . The following lemma of [20] is very useful. (It was given its descriptive name in [5].)

Lemma 11. (No-Cycle Lemma) [20] *If C is r -solvable then there is an r -solution \mathcal{S} for which $G(\mathcal{S})$ is acyclic.*

Another simplifying concept may be assumed by the next lemma. For $1 \leq j \leq t$, let \mathfrak{s}_j be the subsequence of \mathcal{S} moving the pebble p_j from its original position (say a vertex v_j) to the target r (i.e. the subsequence of movements σ_i such that $p_{\sigma_i} = p_j$), and let \mathfrak{s} be the subsequence of \mathcal{S} formed by the remaining pebbling steps. We say that \mathcal{S} is a *tidy* solution if $\mathcal{S} = \mathfrak{s}, \mathfrak{s}_1, \dots, \mathfrak{s}_t$, with each subsequence \mathfrak{s}_j consisting of moving the pebble p_j along an *induced* slide from the vertex v_j to r .

We say that \mathcal{S}' is a *tidy rearrangement* of \mathcal{S} if \mathcal{S}' is a permutation of the pebbling steps of \mathcal{S} and \mathcal{S}' is tidy. Since none of the pebbles p_1, p_2, \dots, p_t is discarded in the solution \mathcal{S} , the pebbling steps of \mathcal{S} can be permuted to create a new t -fold r -solution sequence \mathcal{S}' , namely, $\mathfrak{s}, \mathfrak{s}_1, \dots, \mathfrak{s}_t$. Notice that when \mathcal{S} is *minimum* (i.e., using the fewest pebbling steps), the No-Cycle Lemma 11 implies that each subsequence \mathfrak{s}_j consists of moving the pebble p_j along an *induced* slide from the vertex v_j to r . That is, \mathcal{S}' is tidy. We record this as follows.

Lemma 12. (Tidy Lemma) *If \mathcal{S} is a t -fold r -solution of a configuration C on the graph G , then there is a tidy rearrangement \mathcal{S}' of \mathcal{S} .*

For an r -solution $\sigma = \sigma_1, \dots, \sigma_c$, we define its *cost* to equal $\text{cost}(\sigma) = c + 1$. The idea is that we lose one pebble in each step, plus one more pebble that is placed on r — these pebbles cannot be used in subsequent r -solutions. We say that σ is *cheap* if $\text{cost}(\sigma) \leq 2^d$ (and *super-cheap* if $\text{cost}(\sigma) < 2^d$), and define the parameter $q(G, r)$ to be the minimum number of pebbles m so that every configuration of m pebbles has a cheap r -solution. Of course we always have $q(G, r) \geq \pi(G, r)$. For the particular graph $G = P_n^{(k)}$ with simplicial target r we instead use the notation $q(n, k)$.

We say that a graph G is r -(semi)greedy if every configuration of size at least $\pi(G, r)$ has a (semi)greedy r -solution; that is, every pebbling step in the solution decreases (does not increase) the distance of the moved pebble to r .

Lemma 13. (Cheap Lemma) [3] *Given the graph G with target r let G^* be an r -greedy spanning subgraph of G preserving distances to r . Then any configuration of G of size at least $\pi(G^*, r)$ is cheap; i.e. $q(G, r) \leq \pi(G^*, r)$.*

In particular, if T is a breadth-first-search spanning tree of G , rooted at r , then $q(G, r) \leq \pi(T, r)$. In our case we can choose T to be a caterpillar with main path of length d , so that $\pi(T, r) = 2^d + n - d - 1$; that is, $q(n, k) \leq 2^d + n - d - 1$.

Another useful lemma is the following. The proof given in [3] for the case $t = 1$ extends to all t .

Lemma 14. (Edge Removal Lemma) [3] *Given the graph G with target r , if e is an edge between two neighbors of r then $\pi_t(G, r) = \pi_t(G - e, r)$.*

Let S_i be the slide from v_n to v_1 with interior vertices $v_{i_1}, v_{i_2}, \dots, v_{i_{m_i}}$. Clearly, if $j > j'$, then $i_j \leq i_{j'}$. If S_i and S_{ℓ} are slides such that $i_1 \leq \ell_1$, then we can assume that for every j , $i_j \leq \ell_j$.

Finally, we note a key property of k -connected graphs that follows from Menger's Theorem and Dirac's Fan Lemma (see Exercise 4.2.28 in [25].)

Lemma 15. *Let X and Y be disjoint sets of vertices in a k -connected graph G . For each $x \in X$ and $y \in Y$, let $u(x)$ and $w(y)$ be non-negative integers such that $\sum_{x \in X} u(x) = \sum_{y \in Y} w(y) = k$. Then G has k pairwise internally disjoint (X, Y) -paths such that, for each $x \in X$ and $y \in Y$, $u(x)$ of them start at x and $w(y)$ of them end at y .*

From Lemma 15 we obtain the following simple corollary.

Corollary 16. *Let X and Y be disjoint sets of vertices in a k -connected graph G . For each $x \in X$ and $y \in Y$, let $u(x)$ and $w(y)$ be non-negative integers such that $\sum_{x \in X} u(x) \geq j$ and $\sum_{y \in Y} w(y) \geq j$ for some $j \leq k$. Then G has j pairwise internally disjoint (X, Y) -paths such that, for each $x \in X$ and $y \in Y$, at most $u(x)$ of them start at x and at most $w(y)$ of them end at y .*

Proof. Since G is k -connected it is also j -connected. Choose any set of $u'(x) \leq u(x)$ and $w'(y) \leq w(y)$ such that $\sum_{x \in X} u'(x) = \sum_{y \in Y} w'(y) = j$. Then apply Lemma 15. \square

3.2 Cutting Lemmas

Theorem 17. *Let r be a vertex of a graph G and denote by E_r some set of edges between neighbors of r . Suppose that $G - r - E_r$ has connected components G_1, \dots, G_j for some $j > 1$, and define G'_i to be the subgraph of G induced by $V(G_i) \cup \{r\}$. Then*

$$\pi_t(G, r) = \max_{\sum_{i=1}^j t_i = t+j-1} \sum_{i=1}^j \pi_{t_i}(G'_i, r) - j + 1.$$

Proof. We begin by letting r and E_r be as described in the hypothesis, namely, that $G - r - E_r$ has components G_1, \dots, G_j , for some $j > 1$. By the Edge Removal Lemma 14, $\pi_t(G, r) = \pi_t(G - E_r, r)$.

For any positive t_1, \dots, t_j such that $\sum_{i=1}^j t_i = t + j - 1$ (i.e. $\sum_{i=1}^j (t_i - 1) = t - 1$), there exist t_i -fold r -unsolvable configurations C_i on G'_i of size $\pi_{t_i}(G'_i, r) - 1$. The union $C = \cup_{i=1}^j C_i$ is therefore t -fold

r -unsolvable, showing that $\pi_t(G - E_r, r) \geq 1 + \sum_{i=1}^j (\pi_{t_i}(G'_i, r) - 1) = \sum_{i=1}^j \pi_{t_i}(G'_i, r) - j + 1$. Hence

$$\pi_t(G - E_r, r) \geq \max_{\sum_{i=1}^j t_i = t + j - 1} \sum_{i=1}^j \pi_{t_i}(G'_i, r) - j + 1.$$

Let C be a configuration of maximum size on G that is not t -fold r -solvable, and let C_i denote the restriction of C to G'_i . Without loss of generality $C(r) = 0$, and so $|C| = \sum_{i=1}^j |C_i|$. For each i , C_i is $(t_i - 1)$ -fold but not t_i -fold r -solvable on G'_i ; thus $|C_i| \leq \pi_{t_i}(G'_i, r) - 1$ and $\sum_{i=1}^j (t_i - 1) \leq t - 1$ (i.e. $\sum_{i=1}^j t_i \leq t + j - 1$).

Now the maximality of $|C|$ (and independence of $\{C_i\}$; that is, G'_i and $G'_{i'}$ intersect only on r and so no C_i can affect another $C_{i'}$) implies that $\sum_{i=1}^j (t_i - 1) = t - 1$ and $|C_i| = \pi_{t_i}(G'_i, r) - 1$. Hence $|C| = \sum_{i=1}^j (\pi_{t_i}(G'_i, r) - 1)$ and consequently

$$\pi_t(G - E_r, r) \leq \max_{\sum_{i=1}^j t_i = t + j - 1} \sum_{i=1}^j \pi_{t_i}(G'_i, r) - j + 1.$$

This finishes the proof. □

Because every interior vertex of $P_n^{(k)}$ satisfies the hypothesis of Theorem 17, we obtain the following corollary.

Corollary 18. *Let $r = v_i$ be an interior vertex of $P_n^{(k)}$. By removing the edges $v_h v_j$ between vertices of $N(r)$ with $h < i < j$, we obtain two graphs $P_{n_1}^{(k)}$ and $P_{n_2}^{(k)}$, with $n_1 + n_2 = n + 1$, that are joined at r , which is simplicial in both of them. Then*

$$\pi_t(P_n^{(k)}, r) = \max_{t_1 + t_2 = t + 1} \pi_{t_1}(P_{n_1}^{(k)}, r) + \pi_{t_2}(P_{n_2}^{(k)}, r) - 1.$$

3.3 Chordal Lemmas

Lemma 19. *Let r, v and u be three vertices of a chordal graph G . If v belongs to an induced path between u and r , then any path between u and r contains either v or a neighbor of v . Accordingly, $N[v] \setminus \{r, u\}$ separates r from u .*

Proof. Since v belongs to an induced path between r and u , there is a minimal r - u -separator containing v . Since minimal separators in chordal graphs are cliques, every path between r and u contains either v or a neighbor of v . □

Lemma 20. *Chordal graphs are semi-greedy.*

Proof. Let C be an r -solvable configuration on a chordal graph G . Suppose that \mathcal{S} is a minimum r -solution and that \mathcal{S} is not semi-greedy. Then, after a tidy rearrangement of \mathcal{S} if necessary (by the Tidy Lemma 12), we can assume \mathcal{S} ends by moving a pebble p along an *induced* slide $Q : (u = v_1, v_2, \dots, v_h = r)$ where $\text{dist}(v_i, r) < \text{dist}(v_{i+1}, r)$ for some $i \in \{1, \dots, h-2\}$. By Lemma 19, and the fact that $i+1 < h-1$ (else $\text{dist}(v_i, r) < \text{dist}(v_{i+1}, r) = 1$, which would imply that $v_i = r$), the path $v_{i+1}, v_{i+2}, \dots, v_h = r$ has an interior vertex adjacent to v_i , in contradiction with the fact that Q is an induced slide. \square

Lemma 21. *Let v be a potential vertex of a configuration C on a graph G . Let u be any other vertex which is separated from r by $N[v] \setminus \{r, u\}$. Let C' be the same configuration as C except that the potential vertex v is changed to u ; that is, $C'(v) = C(v) - 2$, $C'(u) = C(u) + 2$, and $C'(x) = C(x)$ otherwise. If C is t -fold r -unsolvable, then C' is t -fold r -unsolvable.*

Proof. Suppose otherwise that C' is t -fold r -solvable and let \mathcal{S} be a minimum t -fold r -solution; by Lemma 20, \mathcal{S} is semi-greedy. We can assume that \mathcal{S} takes a pebble p from u to r . Therefore, by a tidy rearrangement if necessary (by the Tidy Lemma 12), we can also assume that \mathcal{S} ends with a subsequence \mathfrak{s}_p moving the pebble p along an slide Q between u and r . Since $N[v] \setminus \{r, u\}$ separates u from r , there exists an interior vertex of Q , say x , that belongs to $N[v]$. This implies that replacing in \mathcal{S} the subsequence \mathfrak{s}_p by the movement of the pebble p from v to r along the part of the slide Q between x and r , we obtain a semi-greedy t -fold r -solution of the original configuration C , a contradiction. \square

The following theorem is not used in this paper, but is likely to be useful in future work.

Theorem 22. *If G is chordal and C is t -fold r -unsolvable then there is a t -fold r -unsolvable configuration C' with $|C'| = |C|$ and every potential vertex is simplicial.*

Proof. Suppose that C is t -fold r -unsolvable and has a non-simplicial potential vertex v . If v is adjacent to r , the proof is trivial. Otherwise, there exists a simplicial vertex u such that $N[v] \setminus \{r, u\}$ separates u from r . Therefore, the proof follows by Lemma 21. \square

3.4 Pebbling Number for Trees

We will make use of the following well known theorem of Chung [6] on the t -fold pebbling number of a tree with target vertex r , using the notion of a *maximum r -path partition* \mathcal{P} . One can compute such a \mathcal{P} iteratively as follows. Beginning with $H = T$, $W = \{r\}$, and $\mathcal{P} = \emptyset$, we choose a longest path P in H having one endpoint in W . Then we add P to \mathcal{P} , add its vertices to W , remove its edges from H , and repeat. The construction yields $\mathcal{P} = \{P_1, \dots, P_\ell\}$, with subscripts signifying the order of inclusion in \mathcal{P} , and with

$\text{length}(P_i) \geq \text{length}(P_{i+1})$. Note that the number of leaves of T equals $\ell + 1$. In general, we write $\mathcal{P}(T, r)$ for the maximum r -path partition of T .

Theorem 23. [6] *Let T be a tree with maximum r -path partition $\mathcal{P} = \{P_1, \dots, P_\ell\}$, with each P_i having length a_i . Then $\pi_t(T, r) = (t^{2^{a_1}} - 1) + \sum_{i=2}^{\ell} (2^{a_i} - 1) + 1 = t^{2^{a_1}} + \sum_{i=2}^{\ell} 2^{a_i} - \ell + 1$.*

The pebbling number $\pi_t(T)$ is given by choosing r to be a leaf of a longest path of T . We write $\mathcal{P}(T)$ for the maximum path partition of T , equal to $\mathcal{P}(T, r)$ for this choice of r .

3.5 Path Power Lemma

Lemma 24. *For $1 \leq i \leq 2$, let $n_i \geq 1$ and $t_i \geq 1$. If $n_1 + n_2 = n + 1$ and $t_1 + t_2 = t + 1$, then $p_{t_1}(n_1, k) + p_{t_2}(n_2, k) - 1 \leq p_t(n, k)$.*

Proof. Let d_i be the diameter of $P_{n_i}^k$, and d be the diameter of P_n^k . Observe that $d \leq d_1 + d_2 \leq d + 1$. Consider the following three cases.

1. $t_1 \geq t_0(n_1, d_1)$ and $t_2 \geq t_0(n_2, d_2)$.

Suppose, without loss of generality, that $d_1 \geq d_2$. If either $d \leq 2$ or $d_1 = d$ (which implies $d_2 = 1$), the proof is simple, so assume $d \geq 3$ and $d_1 < d$. Because $d_1 + d_2 \geq d$ we have

$$\begin{aligned}
p_{t_1}(n_1, k) + p_{t_2}(n_2, k) - 1 &= l_{t_1}(n_1, k) + l_{t_2}(n_2, k) - 1 \\
&= t_1 2^{d_1} + n_1 - 2 - k(d_1 - 1) + t_2 2^{d_2} + n_2 - 2 - k(d_2 - 1) - 1 \\
&\leq (t_1 2^{d_1} + t_2 2^{d_2}) + (n - 4) - k(d - 2) \\
&= (t_1 2^{d_1} + t_2 2^{d_2}) + (n - 4) - k(d - 1) + k \\
&\leq (t 2^{d_1} + 2^{d_2}) + (n - 4) - k(d - 1) + k \\
&\leq (t 2^{d-1} + 2^1) + (n - 4) - k(d - 1) + k \quad \text{since } d_1 < d \\
&= (t 2^{d-1} + k) + (n - 2) - k(d - 1).
\end{aligned}$$

Note that since $d \geq 3$ we have $d_1 \geq d/2 > 1$. Thus, because $t_1 \geq t_0(n_1, d_1)$, we know that $k \leq t_1(2^{d_1} - 2)/(d_1 - 1) \leq t_1(2^{d_1} - 2) \leq t 2^{d-1}$, and so the above amount is at most $t 2^d + (n - 2) - k(d - 1) = l_t(n, k) \leq p_t(n, k)$.

2. $t_1 \leq t_0(n_1, d_1)$ and $t_2 \leq t_0(n_2, d_2)$.

Here we have

$$\begin{aligned}
p_{t_1}(n_1, k) + p_{t_2}(n_2, k) - 1 &= w_{t_1}(n_1) + w_{t_2}(n_2) - 1 \\
&= 2t_1 + n_1 - 2 + 2t_2 + n_2 - 2 - 1 \\
&= 2(t + 1) + (n + 1) - 5 \\
&= 2t + n - 2 \\
&= w_t(n) \\
&\leq p_t(n, k).
\end{aligned}$$

3. $t_1 > t_0(n_1, d_1)$ and $t_2 < t_0(n_2, d_2)$, or $t_1 < t_0(n_1, d_1)$ and $t_2 > t_0(n_2, d_2)$.

Without loss of generality, we may assume for former, in which case we have $t_1(2^{d_1} - 2) > k(d_1 - 1)$ and $t_2(2^{d_2} - 2) < k(d_2 - 1)$. In particular, each $d_i \geq 2$, and so we will make use of the fact that $\left(\frac{2^j - 2}{j - 1}\right)$ is an increasing function for $j \geq 2$. Since $t_1 > t_0(n_1, d_1)$, this implies that

$$k < t_1 \left(\frac{2^{d_1} - 2}{d_1 - 1} \right) < t \left(\frac{2^d - 2}{d - 1} \right), \quad (1)$$

which means that $p_t(n, k) = l_t(n, k) = t2^d + n - 2 - k(d - 1)$. Now we compute

$$\begin{aligned}
p_t(n, k) - [p_{t_1}(n_1, k) + p_{t_2}(n_2, k) - 1] - 1 &= [t2^d + (n - 2) - k(d - 1)] \\
&\quad - [t_1 2^{d_1} + (n_1 - 2) - k(d_1 - 1) + 2t_2 + (n_2 - 2)] - 1 \\
&= [t2^d - k(d - 1)] - [t_1 2^{d_1} - k(d_1 - 1) + 2t_2] \\
&\geq [t(2^d - 2) - k(d - 1)] - [t_1(2^{d_1} - 2) - k(d_1 - 1)],
\end{aligned}$$

which we will show is positive. Indeed, because of the second inequality in (1) we know that

$$t \left(\frac{2^d - 2}{d - 1} \right) \geq t_1 \left(\frac{2^{d_1} - 2}{d_1 - 1} \right),$$

so that

$$t(2^d - 2) - k(d - 1) \geq t_1(2^{d_1} - 2) \left(\frac{d - 1}{d_1 - 1} \right) - k(d - 1),$$

which we will show is greater than $t_1(2^{d_1} - 2) - k(d_1 - 1)$ as follows, using the first inequality of (1) in

the final step.

$$\begin{aligned}
& \left[t_1(2^{d_1} - 2) \left(\frac{d-1}{d_1-1} \right) - k(d-1) \right] - [t_1(2^{d_1} - 2) - k(d_1 - 1)] \\
&= t_1(2^{d_1} - 2) \left[\frac{d-1}{d_1-1} - 1 \right] - k(d - d_1) \\
&= \left[t_1 \left(\frac{2^{d_1} - 2}{d_1 - 1} \right) - k \right] (d - d_1) \\
&= [t_1(2^{d_1} - 2) - k(d_1 - 1)] \left(\frac{d - d_1}{d_1 - 1} \right) \\
&> 0.
\end{aligned}$$

This completes the proof. □

4 Proof of Theorem 4

In this section we verify the Strong t -Target Conjecture 3 for trees. For a (C, D) -solution σ , define $C[\sigma]$ to be the subconfiguration of C consisting of only those pebbles used by σ , and, for $v \in \dot{D}$, define $\sigma[v]$ to be the pebbling moves of σ consisting of only those pebbles used to solve v .

Proof. The result is trivially true when $s(D) = 1$, which means it is true also for $t = 1$ and for $|V(T)| = 1$, so we assume that $s(D) \geq 2$. We now proceed by induction on any of these parameters. Throughout, we will use the fact that every tree T satisfies $\pi_t(T) \geq \pi_{t-1}(T) + 2$; this is evident from Theorem 23 because $2^{a_1} \geq 2$.

Let $|C| = \pi_t(T) - s(D) + 1$. Then $|C| = \pi(T) + (t-1)2^d - s(D) + 1 \geq \pi(T) + (t-1)(2^d - 1) \geq \pi(T)$. If some $r \in \dot{C} \cap \dot{D}$, then $s(D - r) \geq s(D) - 1$ and so

$$\begin{aligned}
|C - r| &= |C| - 1 \\
&= (\pi_t(T) - s(D) + 1) - 1 \\
&\geq \pi_t(T) - s(D - r) - 1 \\
&\geq \pi_{t-1}(T) - s(D - r) + 1.
\end{aligned}$$

If some $r \in \dot{D}$ has $\text{ecc}(r) < d$, then induction on t implies that C has an r -solution σ with cost at most

$2^{\text{ecc}(r)} \leq 2^d - 1$. Then $s(D - r) \geq s(D) - 1$ and so

$$\begin{aligned} |C - C[\sigma]| &= |C| - \text{cost}(\sigma) \\ &\geq (\pi_t(T) - s(D) + 1) - (2^d - 1) \\ &\geq \pi_{t-1}(T) - s(D - r) + 1. \end{aligned}$$

Similarly, if some $r^2 \in \dot{D}$ (i.e. $D(r) \geq 2$) has $\text{ecc}(r) = d$, then induction implies that C has an r -solution σ with cost at most 2^d . Then $s(D - r) = s(D)$ and so

$$\begin{aligned} |C - C[\sigma]| &= |C| - \text{cost}(\sigma) \\ &\geq (\pi_t(T) - s(D) + 1) - 2^d \\ &= \pi_{t-1}(T) - s(D - r) + 1. \end{aligned}$$

In all three cases, induction on t implies that $C - C[\sigma]$ is $(D - r)$ -solvable, making C r -solvable, and any target can be solved with cost at most $2^{\text{ecc}_T(v)}$.

Therefore we may assume that $\dot{C} \cap \dot{D} = \emptyset$ and that if $r \in \dot{D}$ then r is a singleton leaf with $\text{ecc}(r) = d$. Note that $\pi_t(T) \geq \pi_t(T - v) + 1$ for any leaf v , because each $a_i \geq 1$. Since $t \geq 2$ we may choose distinct targets r and v in \dot{D} . Then

$$\begin{aligned} |C| &= \pi_t(T) - s(D) + 1 \\ &\geq (\pi_{t-1}(T) - s(D) + 1) + 1 \\ &= \pi_{t-1}(T) - s(D - v) + 1, \end{aligned}$$

and so, by induction on t , there is an r -solution σ of cost at most 2^d . Therefore

$$\begin{aligned} |C - C[\sigma]| &\geq (\pi_t(T) - s(D) + 1) - 2^d \\ &= \pi_{t-1}(T) - s(D) + 1 \\ &= \pi_{t-1}(T) - s(D - r) \\ &\geq \pi_{t-1}(T - r) - s(D - r) + 1, \end{aligned}$$

which implies, by induction on t and $|V(T)|$, that $C - C[\sigma]$ is $(D - r)$ -solvable and that any $u \in \dot{D} - r$ has a solution of cost at most $2^{\text{ecc}(u)}$. Combined with the prior r -solution, this proves the theorem. \square

5 Proof of Theorem 5

In this section we verify the Strong t -Target Conjecture 3 for powers of paths. We begin by introducing some important notation.

For pebbling functions F_1 and F_2 , define the function $F_1 \pm F_2$ by $(F_1 \pm F_2)(v) = F_1(v) \pm F_2(v)$, as well as the functions $F_1 \wedge F_2$ and $F_1 \vee F_2$ by $(F_1 \wedge F_2)(v) = \min\{F_1(v), F_2(v)\}$ and $(F_1 \vee F_2)(v) = \max\{F_1(v), F_2(v)\}$. (Thus, if $F = F_1 \wedge F_2$ we have $\dot{F} = \dot{F}_1 \cap \dot{F}_2$, while if $F = F_1 \vee F_2$ we have $\dot{F} = \dot{F}_1 \cup \dot{F}_2$.)

Let F be a pebbling function on $G = P_n^{(k)}$. Denote by $G_{[i,j]}$ the subgraph of G induced by the vertices $\{v_i, v_{i+1}, \dots, v_j\}$, and define $F_{[i,j]}$ to be the restriction of F to $G_{[i,j]}$. Define the *pebbling arrangement* of F , $\mathcal{A}(F) = \langle a_1, \dots, a_m \rangle$, where $m = n + |F|$, to be the sequence

$$\langle v_1, v_{1,1}, \dots, v_{1,F(v_1)}, v_2, v_{2,1}, \dots, v_{2,F(v_2)}, \dots, v_n, v_{n,1}, \dots, v_{n,F(v_n)} \rangle.$$

Also, define the *labeling* of \dot{F} to be the sequence

$$\mathcal{L}(F) = (v_{1,1}, \dots, v_{1,F(v_1)}, v_{2,1}, \dots, v_{2,F(v_2)}, \dots, v_{n,1}, \dots, v_{n,F(v_n)}).$$

For $1 \leq i \leq n$, we write $V_i = \{v_i, v_{i,1}, \dots, v_{i,F(v_i)}\}$. Given a pebbling arrangement $\mathcal{A} = \langle a_1, \dots, a_m \rangle$, define $\mathcal{A}_{\langle i,j \rangle}$ to be $\langle a_i, \dots, a_j \rangle$ if $a_i = v_h$, for some h , and $\langle v_h, a_i, \dots, a_j \rangle$ if $a_i \in V_h - \{v_h\}$, for some h . Define $F_{\langle i,j \rangle}$ to be the pebbling function whose pebbling arrangement $\mathcal{A}(F_{\langle i,j \rangle})$ equals $\mathcal{A}_{\langle i,j \rangle}$.

Given a size t distribution D with $s(D) < n$, let j be the least index with $D(v_j) = 0$ and define the configuration W_D by $W_D(v) = 0$ for all $v \in \dot{D}$, $W_D(v_j) = 2t - 1$, and $W_D(v) = 1$ otherwise. Because $\text{pot}(W_D) = t - 1$, W_D is D -unsolvable. Notice also that $|W_D| = n - s(D) + 2t - 2 = (p_t(n, k) - s(D) + 1) - 1$.

In general, call a configuration C D -small if $\text{pot}(C) = |D| - 1$, and a graph G D -small if some D -unsolvable configuration of maximum size is D -small. We say that G is t -small if it is D -small for all $|D| = t$. In addition, define a maximum-size D -small configuration C to be *canonical* whenever the following holds: if $s(D) < n$ then $C(v) = 0$ for all $v \in \dot{D}$ and $C(v)$ is odd for all $v \notin \dot{D}$; if $s(D) = n$ then, for some v with $D(v) = a = \min_u D(u)$, $C(v) = 2t - a - 1$ and $C(u) = 0$ for all $u \neq v$. A configuration C is called *stacked* if there is some vertex v such that $C(v) > 0$ and $C(u) = 0$ for all $u \neq v$. Observe that the canonical configuration in the $s(D) = n$ case above is stacked. In [7], the following theorem was proven by using the appropriate stacked configuration.

Theorem 25. *If $s(D) = n$ then $\pi(K_n, D) = 2|D| - \min_u D(u)$.*

This yields the following corollary.

Corollary 26. *If $s(D) = n$ then K_n has a canonical D -small configuration.*

Theorem 25 is part of the following, more general “Stacking Theorem” proved in [23] and [24].

Theorem 27. *For every graph G , if $s(D) = n(G)$ then there exists a D -unsolvable configuration of maximum size that is stacked.*

Lemma 28. *If $\pi(P_n^{(k)}, D) \leq n + 2|D| - 2 - s(D) + 1$ and if $s(D) < n$ then $P_n^{(k)}$ has a canonical D -small configuration.*

Proof. Suppose that $s(D) < n$ and let C be a configuration of size $n + 2|D| - 2 - s(D)$ such that $C(v) = 0$ for all $v \in \dot{D}$ and $C(v)$ odd for all $v \notin \dot{D}$. Such a configuration exists because, for $t = |D|$, we can place $(t - 1)$ pairs of pebbles the vertices not in \dot{D} , and then one additional pebble on each such vertex, which amounts to $2(t - 1) + (n - s(D)) = |C|$ pebbles in total. Now, such a C then clearly has $\text{pot}(C) = \text{pot}(C') = t - 1$; that is, C is D -small. Moreover, this implies that $\pi(P_n^{(k)}, D) \geq |C| + 1 = n + 2|D| - 2 - s(D) + 1$, and so $|C| = \pi(P_n^{(k)}, D) - 1$; that is, C is maximal D -unsolvable, and hence canonical. \square

Fact 29. *Suppose that $P_n^{(k)}$ is t -wide, and let $d = \text{diam}(P_n^{(k)})$ and D be a distribution of size t . If $d \geq 2$ then $s(D) < n$. Contrapositively, if $s(D) = n$ then $d = 1$; i.e. $P_n^{(k)} = K_n$.*

Proof. This follows immediately from the inequality $s(D) \leq t \leq k(d - 1)/(2^d - 2) \leq k/2 < n$. \square

We prove Theorem 5 by proving the following stronger result.

Theorem 30. *Let k, n and t be positive integers. Assume that D is a target distribution on $G = P_n^{(k)}$ of size t . Then $\pi(G, D) \leq p_t(n, k) - s(D) + 1$. In addition, if G is t -wide then it is D -small, while if G is strictly t -long then any configuration C on G of size $|C| \geq p_t(n, k) - s(D) + 1$ solves any target $v \in \dot{D}$ with cost at most $2^{\text{ecc}_G(v)}$.*

Proof. For given k, n and t , let D be a target distribution on P_n^k with $t = |D|$ and $s = s(D)$. Observe that the D -unsolvable configuration W_D witnesses that $\pi(P_n^{(k)}, D) \geq p_t(n, k) - s(D) + 1$. Hence, when $P_n^{(k)}$ is t -wide and $s(D) < n$, showing that $\pi(P_n^{(k)}, D) \leq p_t(n, k) - s(D) + 1$ will also imply that $\pi(P_n^{(k)}, D) = p_t(n, k) - s(D) + 1 = |W_D| + 1$, and so the second statement of the theorem follows immediately. Anyway, in the arguments that follow, sometime we first prove the second statement and then we use it to prove the upper bound.

We start using induction on k . The case $k = 1$ is a path, which has been proven in Theorem 4 (the only t -wide path is the 1-wide P_2), so we may assume that $k > 1$.

Now we proceed by induction on n .

5.1 Base Step: $n \leq k + 1$

Here $P_n^{(k)} = K_n$ and $d = 1$, so $p_t(n, k) = 2t + n - 2$. Let C be a pebbling configuration with $|C| \geq 2t + n - 2 - s(D) + 1$ and let $H = C \wedge D$, with $|H| = h$. Define $C' = C - H$ and $D' = D - H$, so that $|C'| = |C| - h$ and $|D'| = |D| - h$. Observe that C' has at least $s(D')$ zeros, so that, by the Potential Lemma 10,

$$\begin{aligned} \text{pot}(C') &\geq \left\lceil \frac{(2t - s(D) - 1 - h) + s(D')}{2} \right\rceil \\ &= (t - h) + \left\lceil \frac{h - 1 + s(D') - s(D)}{2} \right\rceil \\ &\geq (t - h), \end{aligned}$$

since $s(D') \geq s(D) - h$. Hence we can solve the h targets of H identically and then solve the remaining $t - h$ targets of $D - H$ with the $t - h$ potential of $C - H$. This completes the upper bound for the case $n \leq k + 1$. As a consequence, $P_n^{(k)}$ is D -small in this case. In addition, every minimal solution has cost at most $2 = 2^d$.

5.2 Induction Step: $n \geq k + 2$

Here we have $d \geq 2$. We use induction on t .

5.2.1 Base Case: $t = 1$

Here G can be t -long, but never strictly so; thus we will not need to calculate the cost of a solution.

In this case we have $s = 1$. Thus, if $d = 2$ then the result holds by Theorem 3 of [4] because G has a universal vertex. So we will assume that $d \geq 3$. In addition, if $k = 2$, then the result holds by Theorem 3.3 of [2], so we will assume that $k \geq 3$.

Let r be the target vertex and suppose that r is internal. Then $P_n^{(k)} - r$ consists of two components G_1 and G_2 such that, for each i , the subgraph of $P_n^{(k)}$ induced by $V(G_i) \cup \{r\}$ is isomorphic to $P_{n_i}^{(k)}$, with $n_1 + n_2 = n + 1$. Now Corollary 18 states that (since the only solution to $t_1 + t_2 = 2$ is $t_1 = t_2 = 1$)

$$\pi(P_n^{(k)}, D) = \pi_1(P_n^{(k)}, r) = \pi_1(P_{n_1}^{(k)}, r) + \pi_1(P_{n_2}^{(k)}, r) - 1,$$

which we can write as

$$\pi(P_n^{(k)}, D) \leq p_1(n_1, k) + p_1(n_2, k) - 1,$$

by using induction on n . Then Lemma 24 reveals that this is at most $p_1(n, k)$, as required.

Thus we may assume that r is simplicial; i.e. that $D(v_1) = 1$. Let C be of size $p_1(n, k) - 1 + 1$; clearly we can assume $C(v_1) = 0$. Hence there is a big vertex u , which means that there is an empty (v_1, u) -separator; i.e, there are k consecutive empty vertices v_{i+1}, \dots, v_{i+k} , for some $i \geq 1$. Recall that $t_0 = t_0(k, d) = k(d-1)/(2^d - 2)$.

1. **Near Case:** $i = 1$.

Here we have $C(v_2) = \dots = C(v_{k+1}) = 0$. To put a pebble on v_1 , it is enough to put two pebbles on v_{k+1} . Let $G' = G_{[k+1, n]}$, with configuration $C' = C_{[k+1, n]}$. Notice that $G' = P_{n'}^{(k)}$, where $n' = n - k$ and $d' = d - 1$, and that $|C'| = |C|$.

(a) **Subcase:** $1 \leq t_0(k, d)$.

In this case we have $|C| = w_1(n) = n$. If $2 \leq t_0(k, d - 1)$ then, by induction on n , we have $\pi_2(G') = w_2(n') = n - k + 2 < n = |C'|$, and so we can place two pebbles on v_{k+1} .

On the other hand, if $2 > t_0(k, d - 1)$ then $2(2^{d-1} - 2) > k(d - 2)$ and so, by induction on n , we have $\pi_2(G') = 2(2^{d-1}) + (n - k) - k(d - 2) - 2 = n + [(2^d - 2) - k(d - 1)] \leq n = |C'|$; thus we can place two pebbles on v_{k+1} .

We can also conclude that in this case G is D -small.

(b) **Subcase:** $1 > t_0(k, d)$.

In this case we have $|C| = l_1(n, k) = 2^d + n - 2 - k(d - 1)$. Because $k \geq 3$, we know that $k \geq 2$. Thus $-k(d - 2) \geq 2 - k(d - 1)$, which implies that $2(2^{d-1} - 2) - k(d - 2) \geq (2^d - 2) - k(d - 1) > 0$. Hence $2 > t_0(k, d - 1)$.

By induction on n we have $\pi_2(G') = l_2(n', k) = 2(2^{d-1}) + (n - k) - 2 - k(d - 2) = |C'|$, and so we can place two pebbles on v_{k+1} .

2. **Far Case:** $i \geq 2$.

Now we have that the first empty cutset is v_{i+1}, \dots, v_{i+k} , for some $i > 1$. Hence $C(v_i) > 0$ and there is a slide from v_i to v_1 , so we may assume that $C(v_i) = 1$. Therefore, to put a pebble on v_1 , it is sufficient to put one more pebble on v_i from the configuration $C' = C_{[i, n]} - v_i = C_{[i+1, n]}$ on $G' = G_{[i, n]}$. Notice that $G' = P_{n'}^{(k)}$, where $n' = n - i + 1$, and that $|C'| \geq |C| - i + 1$.

The diameters of G and G' are related by

$$k(d' - 1) + b' = n' - 2 = n - i + 1 - 2 = k(d - 1) + b - i + 1,$$

so that $k(d - d') = i + (b' - b) - 1$.

(a) Subcase: $1 \leq t_0(k, d)$.

In this case we have $|C| = w_1(n) = n$. Because, for $d \geq 2$ and fixed k , $t_0(k, d)$ is a strictly decreasing function of d , and $d' \leq d$, we have $1 \leq t_0(k, d')$. Then, by induction on n , we have $\pi_1(G') = w_1(n') = n - i + 1 \leq |C'|$, and so we can place a pebble on v_i .

Again, we can conclude that in this case G is D -small.

(b) Subcase: $1 > t_0(k, d)$.

In this case we have $|C| = l_1(n, k) = 2^d + n - 2 - k(d - 1)$ and $(2^d - 2) > k(d - 1)$. If $1 \leq t_0(k, d')$ then, by induction on n , we have $\pi_2(G') = w_2(n') = n - i + 1 < [2^d + n - 2 - k(d - 1)] - i + 1 \leq |C'|$, and so we can place two pebbles on v_{k+1} .

On the other hand, if $1 > t_0(k, d')$ then $(2^{d'} - 2) > k(d' - 1)$. Because $t_0(k, d)$ is decreasing in d , and $d' \leq d$, we have

$$\begin{aligned} \left(\frac{2^{d'} - 2}{d' - 1} \right) - k &\leq \left(\frac{2^d - 2}{d - 1} \right) - k \\ &\leq \left(\frac{2^d - 2}{d' - 1} \right) - k \left(\frac{d - 1}{d' - 1} \right), \end{aligned}$$

which implies that $(2^{d'} - 2) - k(d' - 1) \leq (2^d - 2) - k(d - 1)$. Hence, by induction on n , we have $\pi_1(G') = 2^{d'} + (n - i + 1) - k(d' - 1) - 2 \leq [2^d + n - 2 - k(d - 1)] - i + 1 \leq |C'|$, and so we can place two pebbles on v_{k+1} , each at cost at most $2^{\text{ecc}_{G'}(v_{k+1})}$, and therefore at total cost at most $2^{\text{ecc}_G(v_{k+1})}$.

This completes the proof for the case $t = 1$.

5.2.2 Inductive Case: $t \geq 2$.

Let C be of size $p_t(n, k) - s(D) + 1$. We will prove C solves D . If $C(x) \wedge D(x) > 0$ for some x , then we use a pebble on x to solve a target on x , and then use $C - x$ to solve $D - x$ by induction on t . This is possible

because $p_t(n, k) > p_{t-1}(n, k) + 2$ for all $d \geq 2$ and all t , and because $s(D - x) \geq s(D) - 1$, so that

$$\begin{aligned} |C - x| &= (p_t(n, k) - s(D) + 1) - 1 \\ &\geq p_{t-1}(n, k) - s(D - x) + 1 . \end{aligned}$$

Hence we may assume that $\dot{C} \cap \dot{D} = \emptyset$ (which in turn implies that $s(D) < n$), and no vertex adjacent to a target vertex is big.

If $C(v_1) = D(v_1) = 0$, then we use induction on n , since

$$\begin{aligned} |C_{[2, n]}| &= p_t(n, k) - s(D) + 1 \\ &\geq [p_t(n - 1, k) - s(D_{[2, n]}) + 1] + 1 . \end{aligned}$$

The same can be said by symmetry for v_n . Hence we may assume that each of v_1 and v_n have either a pebble or a target on it.

We consider three cases regarding the size of t . Recall that $t_0 = t_0(n, k) = k(d - 1)/(2^d - 2)$.

1. **Wide Case:** $t \leq t_0$.

We consider two cases regarding the size of d .

(a) **Subcase:** $d = 2$.

Here we have $t \leq k(d - 1)/(2^d - 2) = k/2$. Let $n = k + 2 + b$ and $|C| = 2t + n - 2 - s + 1$, where $s = s(D)$. Define $M = \{v_{b+2}, \dots, v_{k+1}\}$ and note that every vertex in M is dominating. This means that every pair of potentials can solve any target via pebbling steps through any vertex in M . Because $\dot{C} \cap \dot{D} = \emptyset$, we first observe that $z \geq s$, where z is the number of zeros of C , which, by Lemma 10, implies that

$$\text{pot}(C) \geq \left\lceil \frac{(2t - 2 - s + 1) + s}{2} \right\rceil = t .$$

Since $t \geq 2$ and any pair of potentials solves any root r through any vertex of M , we have a solution at cost $4 = 2^{\text{ecc}(r)}$, because any root has eccentricity 2.

Next, for each $x \in \dot{D}$, define $u(x) = 2D(x)$ and, for each potential vertex y of C , define $w(y) = 2\lfloor C(y)/2 \rfloor$ (i.e. twice the number of potentials at y). Then, since $t \leq k/2$, Corollary 16 (with $j = 2t \leq k$) yields $2t$ internally disjoint (D, B) -paths \mathcal{P} , where B is the set of potentials of

C . Let ρ be the number of targets in \dot{D} that have a path in \mathcal{P} with no zeros on it. Then there are $t - \rho$ targets in \dot{D} , all of whose paths in \mathcal{P} contain at least one zero; that is, there are at least $2(t - \rho)$ zeros of C different from $\text{supp}(D)$. Hence

$$\begin{aligned} \text{pot}(C) &\geq \left\lceil \frac{(2t - 2 - s + 1) + (s + 2t - 2\rho)}{2} \right\rceil \\ &= \left\lceil \frac{4t - 1 - 2\rho}{2} \right\rceil \\ &= 2t - \rho \\ &= \rho + 2(t - \rho). \end{aligned}$$

Thus we can solve the remaining $t - \rho$ targets via pairs of potentials, which solves D . This completes the upper bound for the subcase $d = 2$ of the case $t \leq t_0$. As a consequence, $P_n^{(k)}$ is D -small in this subcase.

- (b) **Subcase: $d \geq 3$.** Notice that, if $t' \leq t$ and $n' \leq n$, then $P_{n'}^{(k)}$ is t' -wide: $d' = \text{diam}(P_{n'}^{(k)}) \leq \text{diam}(P_n^{(k)}) = d$, and so


$$t' \leq t \leq k \left(\frac{d-1}{2^d-2} \right) \leq k \left(\frac{d'-1}{2^{d'}-2} \right),$$

since $d \geq 3$ and the function $(x-1)/(2^x-2)$ is decreasing for $x > 1$. Because $P_{n'}^{(k)}$ is t' -wide, we may assume that $P_{n'}^{(k)}$ is t' -small for all such n' and t' (except for when $n' = n$ and $t' = t$); that is, given any target configuration D' of size t' , among the D' -unsolvable configurations of maximum size there exist at least one with $t' - 1$ potential movements.

Before proving that C solves D , we will prove that $P_n^{(k)}$ is D -small; i.e. we will prove that there exists a pebbling configuration C^* of maximum size among the D -unsolvable configurations s.t. $\text{pot}(C^*) = |D| - 1 = t - 1$.

Write $D = \{v_{i_1}, \dots, v_{i_t}\}$, with $i_j \leq i_{j+1}$ for all $1 \leq j < t$, and let C' be D -unsolvable of maximum size. Moreover, among all such configurations, we may choose C' to have the fewest pebbles in $G_{[1,k]}$; i.e. $c = |C'_{[1,k]}|$ is minimum. Also, since C' is D -unsolvable, we have $|C'| \geq |W_D| \geq w_t(n) - s(D) + 1$, and so, from above, we can assume that $\dot{C}' \cap \dot{D} = \emptyset$, that no big vertex of C' is adjacent to a vertex of D , and that each of v_1 and v_n have either a target or a pebble.

Observe that C' is $(D - v_{i_j})$ -solvable for all $1 \leq j \leq t$. Let $\mathcal{A} = \langle a_1, \dots, a_m \rangle$ be the pebbling arrangement of $C' + D$, with $m = n + |C'| + |D|$. Denote by $C'_{\langle i,j \rangle}$ the portion of C' that is in $\mathcal{A}_{\langle i,j \rangle}$. For each $1 \leq j \leq t$ define ι_j such that $v_{i_j} = a_{\iota_j}$, and define $C'_{L_j} = C'_{\langle 1, \iota_j \rangle}$ and $C'_{R_j} = C'_{\langle \iota_j, n \rangle}$, with



$$\mathcal{A}(C + D) = \langle v_1, v_{1,1}, v_{1,2}, v_2, v_{2,1}, v_{2,2}, v_{2,3}, v_{2,4}, v_3, v_{3,1}, v_{3,2}, v_4, v_{4,1}, v_5, v_{5,1}, v_{5,2}, v_{5,3}, v_6, v_{6,1} \rangle$$

$\begin{array}{c c} j & l_j \\ \hline 1 & 2 \\ 2 & 3 \\ 3 & 10 \\ 4 & 11 \\ 5 & 13 \end{array}$	$D'_{L_1} = \{v_1\}$	$C'_{L_1} = \{\}$
	$D'_{L_2} = \{v_1^2\}$	$C'_{R_2} = \{v_2^4, v_5^3, v_6\}$
	$D'_{R_4} = \{v_3, v_4\}$	$C'_{L_4} = \{v_2^4\}$

Figure 1: Example for $G = P_6^{(2)}$ with $t = 5$, $v_{i_5} = v_4$, and D does not precede C .

$D_{L_j} = \{v_{i_1}, \dots, v_{i_j}\}$ and $D_{R_j} = \{v_{i_j}, \dots, v_{i_t}\}$. See an example in Figure 1.

If some $1 < j < t$ has C'_{L_j} D_{L_j} -unsolvable and C'_{R_j} D_{R_j} -unsolvable then, by induction on t there exist maximum D_{L_j} -unsolvable $C_{L_j}^*$ that is D_{L_j} -small, and maximum D_{R_j} -unsolvable $C_{R_j}^*$ that is D_{R_j} -small. By Corollary 26 and Lemma 28, we may choose both $C_{L_j}^*$ and $C_{R_j}^*$ to be canonical. Because $\dot{C}' \cap \dot{D} = \emptyset$, we have $s(D) < n$, which implies that either $s(D_{L_j}) < i_j$ or $s(D_{R_j}) < n - i_j + 1$. By symmetry, we will assume the former, from which follows that $C_{L_j}^*(v_{i_j}) = 0$. This implies that $\dot{C}_{L_j}^* \cap \dot{C}_{R_j}^* = \emptyset$. Then $C^* = C_{L_j}^* + C_{R_j}^*$ is D -unsolvable and D -small, since $\text{pot}(C^*) = \text{pot}(C_{L_j}^*) + \text{pot}(C_{R_j}^*) = (j - 1) + (t - j) = t - 1$. And it is of maximum size, since $|C^*| = |C_{L_j}^*| + |C_{R_j}^*| \geq |C'_{L_j}| + |C'_{R_j}| = |C'|$ and C' is maximum D -unsolvable (the second equality holds because $C'(v_{i_j}) = 0$, by assumption).

Thus, for all $1 < j < t$, either C'_{L_j} is D_{L_j} -solvable or C'_{R_j} is D_{R_j} -solvable. The same holds true if $j \in \{1, t\}$ and v_{i_j} is not simplicial, although induction is on n instead of t .

If v_{i_1} is not simplicial, then C'_{L_1} must solve D_{L_1} . Let h be the biggest j such that C'_{L_j} solves D_{L_j} . Clearly, $h < t$ and $C'_{R_{h+1}}$ does not solve $D_{R_{h+1}}$, which implies that either $h + 1 = t$ and $v_{i_t} = v_n$ is simplicial, or $C'_{L_{h+1}}$ solves $D_{L_{h+1}}$ in contradiction to the choice of h . We conclude that if v_{i_1} is not simplicial, then v_{i_t} is. Therefore, we may assume that the graph is labeled so that $v_1 \in \dot{D}$.

First suppose that $c = 0$ and set $C'' = C'_{[k+1, n]} = C'$. For $G = P_n^{(k)}$, let $G' = G_{[k+1, n]} \cong P_{n-k}^{(k)}$; then G' has diameter $d' = d - 1$. In this case we use induction on n . We simplify notation somewhat; $D_L = D_{[1, k]}$, $D_R = D_{[k+1, n]}$, $t_L = |D_L|$, and $t_R = |D_R| = t - t_L$. Then set $n' = n - k$ and $t' = t + t_L$.

If G' is t' -wide then define $D' = D_R + 2t_L v_{k+1}$; then $|D'| = t'$. Note that $s(D') = s(D) - s(D_L) + \epsilon$ for some $\epsilon \in \{0, 1\}$. Hence $s(D) - s(D') + 2t_L = s(D_L) - \epsilon + 2t_L \leq 3t_L \leq 3t \leq k$, so that $-s(D) \geq -k + 2t_L - s(D')$. Therefore we have

$$\begin{aligned}
|C''| &= |C'| \\
&\geq n + 2t - 2 - s(D) + 1 \\
&\geq (n - k) + 2(t + t_L) - 2 - s(D') + 1 \\
&= n' + 2t' - 2 - s(D') + 1,
\end{aligned}$$

and so C'' is D' -solvable in G' , which implies that C' solves D in G , a contradiction.

If G' is t' -long then define $D' = D_R + 2D_L^{+k}$, where $D_L^{+k} = \sum_{i=1}^k D(v_i)v_{k+i}$ so that $s(D') = s(D) - s(D_L^{+k} \cap D_R)$. Now we have

$$\begin{aligned}
|C''| &= |C'| \\
&\geq n + 2t - 2 - s(D) + 1 \\
&= k(d - 1) + b + 2t - s(D) + 1 \\
&\geq t(2^d - 2) + b + 2t - s(D) + 1 \\
&\geq t2^d + b - s(D') + 1 \\
&= (t + t_L)2^{d-1} + t_R 2^{d-1} + b - s(D') - s(D_L^{+k} \cap D_R) + 1 \\
&= [t'2^{d'} + b - s(D') + 1] + [t_R 2^{d'} - s(D_L^{+k} \cap D_R)] \\
&\geq [t'2^{d'} + b - s(D') + 1] + [4t_R - 3t_R] \\
&\geq t'2^{d'} + b - s(D') + 1.
\end{aligned}$$

Hence C' is D' -solvable in G' , which implies that C' solves D in G , a contradiction.

These contradictions imply that $c > 0$. We now define j to be minimum so that $v_j \in \dot{C}'$, with $C'' = C - v_j$ and $D' = D + v_j - v_1$. If $D(v_1) = 1$ then $s(D') = s(D)$ and $D'(v_1) = 0$, so set $G' = G - v_1 \cong P_{n'}^{(k)}$, where $n' = n - 1$. Then

$$\begin{aligned}
|C''| &= |C'| - 1 \\
&= (n + 2t - 2 - s(D) + 1) - 1 \\
&= n' + 2t - 2 - s(D') + 1,
\end{aligned}$$

which means, since $P_n^{(k)}$ is t -wide, that C'' is D' -solvable on G' by induction on n . If instead $D(v_1) > 1$ then $s(D') = s(D) + 1$ and $c' = |C''_{[i_{t+1}, k+1]}| = c - 1$, and so

$$\begin{aligned} |C''| &= |C'| - 1 \\ &= (n + 2t - 2 - s(D) + 1) - 1 \\ &= n + 2t - 2 - s(D') + 1, \end{aligned}$$

which means that C'' is D' -solvable on G by minimality of c . In either case, because $j \leq k + 1$, the pebble placed by C'' on v_j , along with the pebble of C' already on v_j , can reach v_1 , which makes C' D -solvable, a contradiction.

This final contradiction implies that C' is D -small, which completes the proof that $P_n^{(k)}$ is D -small.

We complete the proof of this subcase of Theorem 5 as follows. Since $P_n^{(k)}$ is D -small, we have that $\pi(P_n^{(k)}, D) = 1 + |C'|$ s.t. C' is D -small. It is easy to see that if C' is D -small, then $|C'| \leq n - s(D) + 2\text{pot}(C') \leq n - s(D) + 2(t - 1)$, and thus $\pi(P_n^{(k)}, D) \leq 1 + n - s(D) + 2(t - 1) = |C'|$, which implies the result.

2. Intermediate Case: $t_0 < t = \lceil t_0 \rceil$.

(a) Subcase: $d = 2$.

We note that this case follows the same argument as in the $d = 2$ subcase of the wide case $t \leq t_0$; only the size of C is different.

Here we have $t = (k + 1)/2$ for some odd k . Let $n = k + 2 + b$ and $|C| = 4t + b - s + 1$, where $s = s(D)$. Define $M = \{v_{b+2}, \dots, v_{k+1}\}$ and note that every vertex in M is dominating. This means that every pair of potentials can solve any target via pebbling steps through any vertex in M . We first observe that $z \geq s$, which implies that

$$\begin{aligned} \text{pot}(C) &\geq \left\lceil \frac{(4t + b - s + 1 - n) + s}{2} \right\rceil \\ &= t + \left\lceil \frac{2t + b + 1 - n}{2} \right\rceil \\ &= t. \end{aligned}$$

Since $t \geq 2$ and any pair of potentials solves any root r through any vertex of M , we have a

solution at cost $4 = 2^{\text{ecc}(r)}$, because any root has eccentricity 2.

Next, for each $x \in \dot{D}$, define $u(x) = 2D(x)$ and, for each potential vertex y of C , define $w(y) = 2\lfloor C(y)/2 \rfloor$ (i.e. twice the number of potentials at y). Then, since $t = (k+1)/2$, Corollary 16 (with $j = 2t - 1 \leq k$) yields $2t - 1 = k$ internally disjoint (D, B) -paths \mathcal{P} , where B is the set of potentials of C . Let ρ be the number of targets in \dot{D} that have a slide in \mathcal{P} . Then there are $t - \rho$ targets in \dot{D} , all of whose paths in \mathcal{P} contain at least one zero; that is, there are at least $2(t - \rho) - 1$ zeros of C different from $\text{supp}(D)$. Hence

$$\begin{aligned} \text{pot}(C) &\geq \left\lceil \frac{(4t + b - s + 1 - n) + (s + 2t - 2\rho - 1)}{2} \right\rceil \\ &= \left\lceil \frac{4t - 2\rho - 1}{2} \right\rceil \\ &= 2t - \rho \\ &= \rho + 2(t - \rho). \end{aligned}$$

Thus we can solve the remaining $t - \rho$ targets via pairs of potentials, which solves D .

(b) Subcase: $d \geq 3$.

Now we have $p_t(n, k) = l_t(n, k) = t2^d + b = t2^d + (n - 2) - k(d - 1)$. Let $|C| = l_t(n, k) - s(D) + 1$.

Note that, because $t = \lceil t_0 \rceil$, for any $n' = k(d' - 1) + 2 + b$ for some $0 \leq b < k$, $P_{n'}^{(k)}$ is t' -wide for all $t' \leq t$ and $d' = \text{diam}(P_{n'}^{(k)}) \leq \text{diam}(P_n^{(k)}) = d$, provided that either $t' < t$ or $d' < d$:

$$\left(\frac{d - 1}{2^d - 2} \right) \leq \left(\frac{d' - 1}{2^{d'} - 2} \right),$$

and thus

$$t' \leq t = \lceil t_0(k, d) \rceil = \left\lceil k \left(\frac{d - 1}{2^d - 2} \right) \right\rceil \leq \left\lceil k \left(\frac{d' - 1}{2^{d'} - 2} \right) \right\rceil = \lceil t_0(k, d') \rceil.$$

Notice that we get a strict inequality if either $t' < t$ or $d' < d$, which would then imply that $t' \leq t_0(k, d')$; i.e. $P_{n'}^{(k)}$ is t' -wide.

Let $\mathcal{A} = \langle a_1, \dots, a_m \rangle$ be the pebbling arrangement for the pebbling function $C \vee D$. For $1 \leq i < j \leq m$, let $C_{\langle i, j \rangle}$ (resp., $D_{\langle i, j \rangle}$) refer to only those pebbles (resp., targets) of $(C \vee D)_{\langle i, j \rangle}$ that correspond to C (resp., D). Now set $s_{\langle i, j \rangle} = |\text{supp}(D_{\langle i, j \rangle})|$, let h_j be such that $a_j \in V_{h_j}$, and define

$$\Delta(j) = |C_{\langle 1, j \rangle}| - p_{|D_{\langle 1, j \rangle}|}(h_j, k) + s_{\langle 1, j \rangle} - 1$$

Notice that $1 \leq h_j \leq n - b - 1$ implies that

$$\Delta(j) = |C_{\langle 1, j \rangle}| - 2|D_{\langle 1, j \rangle}| - h_j + s_{\langle 1, j \rangle} + 1,$$

because $G_{[1, h_j]}$ is $|D_{\langle 1, j \rangle}|$ -wide in that range. Similarly, the analogous formula holds for $G_{\langle j+1, m \rangle}$ for $b \leq h_{j+1} \leq n$ because it is $|D_{\langle j+1, m \rangle}|$ -wide in that range. The function Δ serves as an indicator: $\Delta(j) \geq 0$ implies that $C_{\langle 1, j \rangle}$ solves $D_{\langle 1, j \rangle}$. If $\Delta(j)$ is too large, $|C_{\langle j+1, m \rangle}|$ may be too small to solve $D_{\langle j+1, m \rangle}$. However, if $\Delta(j) = \epsilon \in \{0, 1\}$ for some j for which both $G_{\langle 1, j \rangle}$ is $|D_{\langle 1, j \rangle}|$ -wide and $G_{\langle j+1, m \rangle}$ is $|D_{\langle j+1, m \rangle}|$ -wide, then we can show that $C_{\langle j+1, m \rangle}$ solves $D_{\langle j+1, m \rangle}$. Indeed,

$$\begin{aligned} |C_{\langle j+1, m \rangle}| &= |C| - |C_{\langle 1, j \rangle}| \\ &= (t2^d + b - s(D) + 1) - (2|D_{\langle 1, j \rangle}| + h_j - s_{\langle 1, j \rangle} - 1 + \epsilon) \\ &\geq (2t + n - 2 - s(D) + 2) - (2|D_{\langle 1, j \rangle}| + h_j - s_{\langle 1, j \rangle} - 1 + \epsilon) \\ &= 2(t - |D_{\langle 1, j \rangle}|) + (n - h_j) - (s(D) - s_{\langle 1, j \rangle}) + (1 - \epsilon) \\ &\geq 2|D_{\langle j+1, m \rangle}| + (n - h_j + 1) - 2 - s_{\langle j+1, m \rangle} + 1 \\ &\geq p_{|D_{\langle j+1, m \rangle}|}(G_{\langle j+1, m \rangle}) - s_{\langle j+1, m \rangle} + 1 \\ &\geq \pi(P_{n-h_{j-1}+1}^{(k)}, D_{\langle j+1, m \rangle}). \end{aligned}$$

Therefore, we aim to show that such a j exists. Let j^* be maximum such that $G_{\langle 1, j^* \rangle}$ is $|D_{\langle 1, j^* \rangle}|$ -wide. Then $G_{\langle 1, j \rangle}$ is $|D_{\langle 1, j \rangle}|$ -wide for all $1 \leq j \leq j^*$. Thus

$$\Delta(j) - \Delta(j-1) = \begin{cases} -1 & \text{for } a_j = v_h, \text{ for some } h, \\ +1 & \text{for } a_j \in \dot{C}, \\ -1 & \text{for } a_j \in \dot{D} \text{ and } a_j = v_{h,1}, \text{ for some } h, \text{ and} \\ -2 & \text{for } a_j \in \dot{D} \text{ and } a_j = v_{h,\ell}, \text{ for some } h \text{ and some } \ell > 1. \end{cases} \quad (2)$$

Moreover, for $j > j^*$, even in the cases for which $G_{\langle 1, j \rangle}$ is $|D_{\langle 1, j \rangle}|$ -long, one can see that

$$\Delta(j) - \Delta(j-1) > 0 \text{ if and only if } a_j \in \dot{C}, \text{ in which case } \Delta(j) - \Delta(j-1) = 1. \quad (3)$$

Indeed, assume that $G_{\langle 1, j \rangle}$ is t -long.

- Consider when $|D_{\langle 1, j-1 \rangle}| = t$. If $G_{\langle 1, j-1 \rangle}$ is t -long then $a_j \in V(G) \cup \dot{C}$ since $D_{\langle j, m \rangle}$ is empty.

In this case,

$$\Delta(j) - \Delta(j-1) = \begin{cases} -1 & \text{for } a_j = v_h, \text{ for some } h, \text{ and} \\ +1 & \text{for } a_j \in \dot{C}. \end{cases}$$

If $G_{\langle 1, j-1 \rangle}$ is t -wide then $a_j \in V_{(d-1)k+2}$, and so $\Delta(j) - \Delta(j-1) \leq (d-1)k - t(2^d - 2) - 1 \leq -2$.

- Now consider when $|D_{\langle 1, j-1 \rangle}| < t$. Then $G_{\langle 1, j-1 \rangle}$ is $(t-1)$ -wide, and so $a_j \in \dot{D}$. In this case,

$$\Delta(j) - \Delta(j-1) \leq \begin{cases} -2 & \text{for } a_j \in \dot{D} \text{ and } a_j = v_{h,1}, \text{ for some } h, \text{ and} \\ -3 & \text{for } a_j \in \dot{D} \text{ and } a_j = v_{h,\ell}, \text{ for some } h \text{ and some } \ell > 1. \end{cases}$$

This proves (3).

Suppose that $D(v_1) > 0$ and $D(v_n) > 0$. Then $\Delta(2) = -1$ and $\Delta(m) = 0$. Let j' be minimum such that $\Delta(j') = 0$, with $a_j \in V_h$. Because of Equations (2) and (3), we must have both $a_{j'-1}$ and $a_{j'}$ in \dot{C} . Indeed, $\Delta(j') = 0$ implies that $\Delta(j'-1) = -1$ and $\Delta(j'-2) = -2$ by the definition of j' . This means that $\Delta(j') - \Delta(j'-1) = 1$ and $\Delta(j'-1) - \Delta(j'-2) = 1$, so that $a_{j'}$ and $a_{j'-1}$ are both pebbles; i.e. v_h is big. Since no big vertex is adjacent to a target, we have $k+1 < h < n-k$, which proves the existences of the desired j .

Now suppose that $D(v_1) = 0$ or $D(v_n) = 0$. By symmetry we will assume that $D(v_1) = 0$. Suppose that $v_h \in \dot{D}$ for some $1 < h \leq k+1$. Because no big vertex is adjacent to a target vertex, $C(v_i) \leq 1$ for all $i < h$; thus $|C_{[1, h-1]}| \leq h-1$. Therefore

$$\begin{aligned} |C_{[h, n]}| &= |C| - |C_{[1, h-1]}| \\ &\geq (t2^d + n - 2 - k(d-1) - s(D) + 1) - (h-1) \\ &= t2^d + (n - h + 1) - 2 - k(d-1) - s(D) + 1. \end{aligned}$$

If $h \leq b$ then P_{n-h+1} is t -long, and so this value equals $p_t(P_{n-h+1}^{(k)}) - s(D) + 1$, implying that $C_{[h, n]}$ solves D by induction on n . If $b < h \leq k+1$ then P_{n-h+1} is t -wide, and so this value is at least $(n-h+1) + 2t - 2 - s(D) + 1 = p_t(P_{n-h+1}^{(k)}) - s(D) + 1$, implying that $C_{[h, n]}$ solves D by induction on n .

Thus we may assume that $D_{[1, k+1]}$ is empty. Notice that the above arithmetic also proves that if h is the minimum index such that $v_h \in \dot{D}$, then $|C_{[1, h-1]}| \geq h$: if $|C_{[1, h-1]}| \leq h-1$ then remove $G_{[1, h-1]}$; then $C_{[h, n]}$ solves D by induction on n , because $G_{[h, n]}$ is t -wide, since $h > k+1$. We will make use of this below.

Let j_0 be minimum so that $a_{j_0} \in \dot{D}$; thus $a_{j_0} \in V_h$. Because $|C_{\langle 1, j_0 \rangle}| = |C_{[1, h-1]}| \geq h \geq$

$\pi(G_{\langle 1, j_0 \rangle}, v_h)$ ($G_{\langle 1, j_0 \rangle}$ is 1-wide because $1 < t$), we have $\Delta(j_0) \geq 0$. If $\Delta(j_0) \in \{0, 1\}$ then we are done; thus we will assume that $\Delta(j_0) > 1$. Now let j^* be maximum so that $a_{j^*} \in \dot{D}$ and let h^* be such that $a_{j^*} \in V_{h^*}$. Of course, if $\Delta(j^*) \geq 0$ then we are done, so assume otherwise; i.e. $\Delta(j^*) < 0$.

By the definition of j^* , we have that $G_{\langle 1, j \rangle}$ is $|D_{\langle 1, j \rangle}|$ -wide for all $1 \leq j < j^*$. Thus Equation (2) applies. Because Δ never decreases by more than 2, this implies the existence of some $j_0 \leq j' < j^*$ with $\Delta(j') \in \{0, 1\}$, completing the proof in this case.

3. Long Case: $t > \lceil t_0 \rceil$.

Here we have $p_t(n, k) = l_t(n, k) = t2^d + (n - 2) - k(d - 1)$. Let $|C| = l_t(n, k) - s(D) + 1$. In this case, $t - 1 \geq \lceil t_0 \rceil$, and so $(t - 1)(2^d - 2) \geq k(d - 1)$, which implies that $t2^d - k(d - 1) \geq 2^d - 2 + 2t \geq 2^d - d + 1$ since $2t + d \geq 3$. Hence

$$t2^d + n - 2 - k(d - 1) \geq 2^d + n - d - 1. \quad (4)$$

Let T_i be a breadth-first-search spanning tree of G , rooted at v_i (with T_1 isomorphic to a caterpillar with main path of length d), then $\pi(T_i, v_i) \leq \pi(T_1, v_1) = 2^d + n - d - 1$. Then inequality (4) implies that $|C| \geq \pi(T_1, v_1) - s(D) + 1 \geq \pi(T_i, v_i) - s(D) + 1$ and so, by Theorem 4, C solves any target $v_i \in \dot{D}$ along T_i with cost at most $2^{\text{ecc}_{T_i}(v_i)}$.

If some $x \in \dot{D}$ has $D(x) \geq 2$ or has $\text{ecc}_{T_i}(x) < d$, then

$$\begin{aligned} |C - C[\sigma_x]| &\geq (p_t(n, k) - s(D) + 1) - 2^{\text{ecc}_{T_i}(x)} \\ &= (p_{t-1}(n, k) - s(D - x) + 1) + (2^d - 2^{\text{ecc}_{T_i}(x)}) + (s(D - x) - s(D)) \\ &\geq \pi(G, D - x), \end{aligned} \quad (5)$$

and so $C - \sigma_x$ solves $D - x$.

Hence D consists of singleton targets only, each with eccentricity d . That is, $D = D_{[1, b+1]} \cup D_{[n-b, n]}$; i.e. $\dot{D} \subseteq V_{[1, b+1]} \cup V_{[n-b, n]}$. Without loss of generality, we may assume that $\dot{D}_{[1, b+1]} \neq \emptyset$. In addition, if $\text{cost}(\sigma_{v_i}) = 2^d$ for $i \in [1, b + 1]$ then every potential vertex v_j in $C[\sigma_{v_i}]$ is at distance d from v_i ; that is, $j \in [n - b + i - 1, n]$. Since no big vertex is adjacent to a target, we then have $\dot{D}_{[n-b, n]} = \emptyset$; i.e. $D = D_{[1, b+1]}$. Furthermore, if any length- d path P from a *huge* ($C(v) \geq 2^{\text{dist}(v, r)}$ for some $r \in \dot{D}$) vertex v to some target r contains an interior pebble, then there is a super-cheap r -solution from v along P . The calculations of Inequality (5) show that the existence of a cheap solution yields a t -fold solution by induction on t .

Now, if v_j is big, then $j \in [n - b + i - 1, n]$, so define C' by $C'(v_j) = C(v_j) - 2$, $C'(v_n) = C(v_n) + 2$, and $C'(v) = C(v)$ otherwise. Then any (C', D) -solution can be converted to a (C, D) -solution because $N(v_n) \subseteq N[v_j]$. Thus, if C is not D -solvable, then neither is C' . In other words, we may assume that the only big vertex of C is v_n .

This implies that, for every $v_i \in \dot{D}$, the rail $R_i = (v_i, v_{i+k}, \dots, v_{i+(d-1)k})$ has no pebbles. Let j be minimum such that $v_j \in \dot{D}$. Now define the graph $G' = G - \{R_i\}_{v_i \in \dot{D} - v_j}$, with corresponding configuration $C' = C_{G'}$, and notice that $G' \cong P_{n'}^{(k')}$, where $n' = n - (t-1)d$ and $k' = k - (t-1)$. In addition, $b' = b - (t-1)$ and $d' = d$. By induction on n , we have that

$$\begin{aligned} |C'| &= |C| \\ &= (t2^d + b) - s(D) + 1 \\ &= t2^{d'} + b' \\ &= p_t(n', k'). \end{aligned}$$

which implies that C' is t -fold v_j -solvable on G' . Equivalently, C is t -fold v_j -solvable on G , using only the vertices of G' . Because $G_{[1, k+1]}$ is a clique, any step $v_{i'} \mapsto v_j$ can be converted instead to $v_{i'} \mapsto v_{j'}$ for any $v_{j'} \in \dot{D}$, thus solving D .

This completes the proof. □

6 Proof of Theorem 6

Lemma 31. *If r is a simplicial vertex of $P_n^{(k)}$, then $\pi_t(P_n^{(k)}, r) \geq p_t(n, k)$.*

Proof. Let r be the simplicial vertex v_1 of $P_n^{(k)}$. For $t \geq 1$ we define the configurations $W = W_{t,n}$ and $L = L_{t,n,k}$ on $P_n^{(k)}$ with target r as follows.

$$W(v_i) = \begin{cases} 0, & \text{for } i = 1; \\ 1, & \text{for } 2 \leq i \leq n-1; \\ 2t-1, & \text{for } i = n. \end{cases} \quad L(v_i) = \begin{cases} 0, & \text{for } 1 \leq i \leq (d-1)k+1; \\ 1, & \text{for } (d-1)k+1 < i \leq n-1; \\ t2^d-1, & \text{for } i = n. \end{cases}$$

Notice that $|W| = w_t(n) - 1$ and $|L| = l_t(n, k) - 1$. Let V_i denote the vertices at distance i from v_1 .

Since $\text{pot}(W) = t - 1$, Fact 9 shows that W is t -fold r -unsolvable.

Since $\text{pot}(L) = t2^{d-1} - 1$, L can only place at most $t2^{d-1} - 1$ pebbles on V_{d-1} . If $d = 1$ then $\text{pot}(L) = t - 1$, so Fact 9 shows that L is t -fold r -unsolvable. If $d > 1$ then, by induction on d , L can only place at most $t2^{d-j} - 1$ pebbles on V_{d-j} for each $1 \leq j \leq d$. Thus L can only place at most $t - 1$ pebbles on r , showing that L is t -fold r -unsolvable. \square

Proof of Theorem 6. Lemma 31 provides the lower bound. Theorem 5 provides the upper bound. \square

7 Proof of Corollary 8

Proof. The values for $n \leq 8$ are easy to check, so we assume that $n \geq 9$. We know that k and d must be related to n by $k(d-1) = n - 2 - b$, for some $0 \leq b < k$; that is,

$$n - 2 - k < k(d-1) \leq n - 2. \quad (6)$$

Then $e_\pi(P_n)$ is the least value of k for which this relation holds with

$$k \geq (2^d - 2)/(d-1). \quad (7)$$

Given $n \geq 9$, define $\kappa = M(n) + 1 = \lfloor (n-2)/(\lceil \lg n \rceil - 2) \rfloor + 1$ and $\delta = \lfloor (n-2)/\kappa \rfloor + 1$, so that when $k = \kappa$ we have $d = \delta$. Thus the upper bound follows by showing that (7) also holds with these values of k and d .

We first observe that the definition of κ implies that

$$n - 2 = (\kappa - 1)(\lceil \lg n \rceil - 2) + \ell, \quad (8)$$

for some $0 \leq \ell < \lceil \lg n \rceil - 2$. Define $\bar{\ell} = (\lceil \lg n \rceil - 2) - \ell$; then

$$\kappa = \frac{n - 2 - \ell}{\lceil \lg n \rceil - 2} + 1 = \frac{(n - 2) + \bar{\ell}}{\lceil \lg n \rceil - 2},$$

so that

$$\begin{aligned} \frac{n - 2}{\kappa} &= \frac{(n - 2)(\lceil \lg n \rceil - 2)}{(n - 2) + \bar{\ell}} \\ &= (\lceil \lg n \rceil - 2) - \frac{\bar{\ell}(\lceil \lg n \rceil - 2)}{(n - 2) + \bar{\ell}}, \end{aligned}$$

and

$$\lceil \lg n \rceil - 3 \leq (\lceil \lg n \rceil - 2) - \frac{(\lceil \lg n \rceil - 2)^2}{n} \leq (\lceil \lg n \rceil - 2) - \frac{\bar{\ell}(\lceil \lg n \rceil - 2)}{(n-2) + \bar{\ell}} < \lceil \lg n \rceil - 2.$$

Hence $\delta - 1 = \lfloor (n-2)/\kappa \rfloor = \lceil \lg n \rceil - 3$. This yields $2^\delta = 2^{\lceil \lg n \rceil - 2} \leq 2^{\lg n - 1} = n/2$. Therefore, when $n \geq 9$ we have

$$\begin{aligned} \kappa \left\lfloor \frac{n-2}{\kappa} \right\rfloor &= \left(\left\lfloor \frac{n-2}{\lceil \lg n \rceil - 2} \right\rfloor + 1 \right) (\lceil \lg n \rceil - 3) \\ &> \left(\frac{n-2}{\lceil \lg n \rceil - 2} \right) (\lceil \lg n \rceil - 3) \\ &\geq (n-4)/2 \\ &\geq 2^\delta - 2, \end{aligned}$$

finishing the proof of the upper bound.

For the lower bound, the values for $n \leq 2^{10}$ can be checked by computer, so we assume that $n > 2^{10}$. Given n , we define $\lambda = \lceil \lg n \rceil - 2 \geq 9$, $N = n - 2^{\lambda+1}$, $M = \lfloor (n-2)/\lambda \rfloor$, and $L = (n-2) \bmod \lambda$. Then $n-2 = M\lambda + L$, $0 < N \leq 2^{\lambda+1}$ and $0 \leq L < \lambda$. Furthermore, we define $m = \lfloor (n-2)/\lambda^2 \rfloor$, and $\ell = (n-2) \bmod \lambda^2$, so that $n-2 = m\lambda^2 + \ell$, with $0 \leq \ell < \lambda^2$. Finally, define $\kappa' = M - m - 1$ and $\delta' = \lfloor (n-2)/\kappa' \rfloor + 1$, so that (6) holds when $k = \kappa'$ and $d = \delta'$. Thus the lower bound follows by showing that (7) fails with these values of k and d .

We begin by rewriting κ' as

$$\begin{aligned} \kappa' &= \frac{2^{\lambda+1} - 2 - L + N}{\lambda} - \frac{2^{\lambda+1} - 2 - \ell + N}{\lambda^2} - 1 \\ &= \frac{\lambda(2^{\lambda+1} - 2 - L + N) - (2^{\lambda+1} - 2 - \ell + N) - \lambda^2}{\lambda^2}. \end{aligned}$$

Then we show that $\delta' - 1 = \lambda + 1$ as follows. For the upper bound we use the fact that $\ell - \lambda L \geq -(\lambda-1)^2$, while for the lower bound we use $\ell - \lambda L \leq \lambda(\lambda-1)$. Additionally, we use the following inequality, which can be easily checked to hold for $\lambda \geq 6$:

$$\frac{\lambda^2(2^{\lambda+1} - 2)}{(\lambda-1)(2^{\lambda+1} - 2) - (\lambda-1)^2 - \lambda^2} < \lambda + 2. \quad (9)$$

Then

$$\begin{aligned}
\delta' - 1 &= \left\lfloor \frac{\lambda^2(2^{\lambda+1} - 2 + N)}{\lambda(2^{\lambda+1} - 2 - L + N) - (2^{\lambda+1} - 2 - \ell + N) - \lambda^2} \right\rfloor \\
&= \left\lfloor \frac{\lambda^2(2^{\lambda+1} - 2 + N)}{(\lambda - 1)(2^{\lambda+1} - 2 + N) + (\ell - \lambda L) - \lambda^2} \right\rfloor \\
&\leq \left\lfloor \frac{\lambda^2(2^{\lambda+1} - 2 + N)}{(\lambda - 1)(2^{\lambda+1} - 2 + N) - (\lambda - 1)^2 - \lambda^2} \right\rfloor \\
&\leq \left\lfloor \frac{\lambda^2(2^{\lambda+1} - 2)}{(\lambda - 1)(2^{\lambda+1} - 2) - (\lambda - 1)^2 - \lambda^2} \right\rfloor \\
&\leq \lambda + 1,
\end{aligned}$$

the last inequality because of (9). Also

$$\begin{aligned}
\delta' - 1 &= \left\lfloor \frac{\lambda^2(2^{\lambda+1} - 2 + N)}{(\lambda - 1)(2^{\lambda+1} - 2 + N) + (\ell - \lambda L) - \lambda^2} \right\rfloor \\
&\geq \left\lfloor \frac{\lambda^2(2^{\lambda+1} - 2 + N)}{(\lambda - 1)(2^{\lambda+1} - 2 + N) - \lambda} \right\rfloor \\
&\geq \left\lfloor \frac{\lambda^2}{(\lambda - 1)} \right\rfloor \\
&\geq \left\lfloor \frac{\lambda^2 - 1}{(\lambda - 1)} \right\rfloor \\
&= \lambda + 1.
\end{aligned}$$

Now we show that inequality (7) fails when $k = \kappa$ and $d = \delta$. To do this, recall from above that $N \leq 2^{\lambda+1}$ and $\ell - \lambda L \leq \lambda(\lambda - 1)$. Also, we note that $2\lambda^3 + \lambda^2 - \lambda + 2 < 2^{\lambda+2}$ for $\lambda \geq 9$. Then

$$\begin{aligned}
\kappa'(\delta' - 1) &= \left(\frac{\lambda(2^{\lambda+1} - 2 - L + N) - (2^{\lambda+1} - 2 - \ell + N) - \lambda^2}{\lambda^2} \right) (\lambda + 1) \\
&= \frac{(\lambda^2 - 1)(2^{\lambda+1} - 2 + N) + (\lambda + 1)(\ell - \lambda L + \lambda^2)}{\lambda^2} \\
&\leq \frac{(\lambda^2 - 1)(2^{\lambda+2} - 2) + (\lambda + 1)(\lambda(\lambda - 1) + \lambda^2)}{\lambda^2} \\
&= 2^{\lambda+2} - 2 - \frac{(2^{\lambda+2} - 2) - (2\lambda^3 + \lambda^2 - \lambda)}{\lambda^2} \\
&= 2^{\lambda+2} - 2 - \frac{2^{\lambda+2} - (2\lambda^3 + \lambda^2 - \lambda + 2)}{\lambda^2} \\
&< 2^{\lambda+2} - 2 \\
&= 2^{\delta'} - 2.
\end{aligned}$$

This finishes the proof of the lower bound and, hence, the corollary. \square

8 Comments

We observe the following simple corollary.

Corollary 32. *Suppose that $P_n^{(k)}$ is t -wide, $d = \text{diam}(P_n^{(k)})$, and D is a distribution of size t . Then*

$$\pi(P_n^{(k)}, D) = \begin{cases} 2t - \min_v D(v) & \text{if } s(D) = n, \text{ and} \\ n + 2t - 1 - s(D) & \text{if } s(D) < n. \end{cases}$$

Proof. The $s(D) = n$ case follows from Fact 29 and Theorem 25, while the $s(D) < n$ case follows from Theorem 30 and Lemma 28. \square

We finish with a few comments, conjectures, and open problems that are driven by this work.

1. It would be interesting to calculate a precise formula for $\pi(P_n^{(k)}, D)$ when $P_n^{(k)}$ is t -long.
2. Does the above proof yields an efficient algorithm for D -solving pebbling configurations of size at least $\pi_t(P_n^{(k)}) - s(D) + 1$?
3. Is there a formula for $\pi(T, D)$ for any distribution D on any tree T (akin to the $|D| = 1$ case)? If so, what is it and how is it constructed?
4. We conjecture that the pebbling number of a chordal graph is always witnessed at a simplicial vertex; that is, if G is chordal then there is some simplicial r such that $\pi(G) = \pi(G, r)$.
5. What is the right generalization of Theorem 22 for a general target D on a chordal graph?
6. Note that $\pi(P_{d+1} \square K_k) = k2^d$ was proved in [6]. For $n = k(d+1)$, we have $P_{d+1} \square K_k \subset P_n^{(k)} \subset P_d \boxtimes K_k$, so that

$$\pi_t(P_{d+1} \boxtimes K_k) \leq \pi_t(P_n^{(k)}) \leq \pi_t(P_{d+1} \square K_k) \leq t\pi(P_{d+1} \square K_k) = tk2^d.$$

One can view $P_{d+1} \boxtimes K_k$ as a “blow-up” of the path P_d by cliques K_k : each vertex of P_d is replaced by a clique of size k , and vertices from consecutive cliques are adjacent. More generally we could blow up the path vertices by different amounts, using the notation $K_{k_0} \boxtimes K_{k_2} \boxtimes \cdots \boxtimes K_{k_d}$, which is what is studied by Sieben in [22]. In this context, for $n = k(d-1) + b + 2$ and $G = K_1 \boxtimes K_k \boxtimes \cdots \boxtimes K_k \boxtimes K_{b+1}$, we have $P_n^{(k)} \subset G$, so that $\pi(P_n^{(k)}) = \pi(G)$, since the arguments of Lemma 31 work on G as well.

7. Might some of the methods developed here be useful in lowering the upper bound of $k(d) \leq 2^{2d+3}$ from [8] mentioned in the Introduction?

References

- [1] L. Alcón, M. Gutierrez, and G. Hurlbert, *Pebbling in split graphs*, SIAM J. Discrete Math. **28**, no. 3 (2014), 1449–1466.
- [2] L. Alcón, M. Gutierrez, and G. Hurlbert, *Pebbling in 2-paths*, Elec. Notes Discrete Math., **150** (2015), 145–150.
- [3] L. Alcón, M. Gutierrez, and G. Hurlbert, *Pebbling in semi-2-trees*, Discrete Math. **340** (2017), 1467–1480.
- [4] L. Alcón, M. Gutierrez, and G. Hurlbert, *t-Pebbling in k-connected graphs with a universal vertex*, Matemática Contemporânea **46** (2019), 38–46.
- [5] D. Bunde, E. Chambers, D. Cranston, K. Milans, and D. West, *Pebbling and optimally pebbling in graphs*, J. Graph Th. **57** (2008), 215–238.
- [6] F.R.K. Chung, *Pebbling in hypercubes*, SIAM J. Discrete Math. **2** (1989), no. 4, 467–472.
- [7] B. Crull, T. Cundiff, P. Feltman, G. Hurlbert, L. Pudwell, Z. Szaniszló, and Z. Tuza, *The cover pebbling number of graphs*, Discrete Math. **296** (2005), no. 1, 15–23.
- [8] A. Czygrinow, H. Kierstead, G. Hurlbert, and W. T. Trotter, *A note on graph pebbling*, Graphs and Combin. **18** (2002), 219–225.
- [9] D. Herscovici, B. Hester, and G. Hurlbert, *t-Pebbling and extensions*, Graphs and Combin. **29** (2013), no. 4, 955–975.
- [10] M. Hoffmann, J. Matoušek, Y. Okamoto, and P. Zumstein, *The t-pebbling number is eventually linear in t*, Electron. J. Combin. **18** (2011), no. 1, Paper 153, 4 pp.
- [11] G. Hurlbert, *General graph pebbling*, Discrete Appl. Math. **161** (2013), 1221–1231.
- [12] G. Hurlbert, *Graph Pebbling*, in Modern Methods in Combinatorics 2nd Puntana School of Combinatorics, D. Jaume and S. Eliahou, eds., Centre International de Mathématiques Pures et Appliquées, 2013.
- [13] G. Hurlbert, *Graph Pebbling*, in Handbook of Graph Theory (2nd ed.), Discrete Mathematics and its Applications, J. Gross, J. Yellen, and P. Zhang, eds., CRC Press, Boca Raton, 2014.
- [14] G. Hurlbert, *The weight function lemma for graph pebbling*, J. Combin. Opt. **34**, no. 2 (2017), 343–361.

- [15] G. Hurlbert and E. Seddiq, *On the target pebbling conjecture*, arXiv:2011.10623.
- [16] J.Y. Kim, *Pebbling exponents of graphs*, J. Natural Sci. Catholic Univ. of Daegu **2**, no. 1 (2004), 1–7.
- [17] J.Y. Kim and S.A. Kim; *The pebbling exponent of paths*, Honam Math. J. **32**, no. 4 (2010), 769–776.
- [18] A. Lourdusamy and T. Mathivanan, *The t -pebbling number of squares of cycles*, J. Prime Res. Math. **11** (2015), 61–76.
- [19] K. Milans and B. Clark, *The complexity of graph pebbling*, SIAM J. Discrete Math. **20** (2006), 769–798.
- [20] D. Moews, *Pebbling graphs*, J. Combin. Theory Ser. B **55**, no. 2 (1992), 244–252.
- [21] L. Pachter, H. Snevily, and B. Voxman, *On pebbling graphs*, Congr. Numer. **107** (1995), 65–80.
- [22] N. Sieben, *t -Pebbling in a path of complete graphs*, preprint.
- [23] J. Sjöstrand, *The cover pebbling number of graphs*, Electron. J. Combin. **12** (2005), Note 22, 5 pp.
- [24] A. Vuong and I. Wyckoff, *Conditions for weighted cover pebbling of graphs*, arXiv:math/0410410 (2004).
- [25] D. West, *Introduction to Graph Theory* (2nd ed.), Pearson, London, 2000.