# Section 4.2 The Mean Value Theorem

The Mean Value theorem is a significant theoretical result. Although you will not use it often, it lays the foundation for Chapter 5, and is used in proofs of many fundamental results. We first address a prelimanary result called Rolle's Theorem

**Rolles Theorem** Suppose a function $f(x)$ is continuous on an interval $[a,b]$ and differentiable on the interval $(a,b)$. (That is, $f'(x)$ exists and is defined at every number $x \in (a,b)$.) If $f(a)=f(b)$ then there exists at least one number $c \in (a,b)$ for whic $f'(c)=0$.




← Rolles Theorem guarantees there is c number with $f'(c)=0$, as illustrated. Note that there may be several such c, as is the case here.

The text gives a careful proof of Rolle's theorem, which you should read. But notice the Theorem is very intuitive. Move a horizontal line down (or up) until it hits a point on the graph of $y=f(x)$. The point of first contact has an x-coordinate c for which $f'(c)=0$, as illustrated here




From Rolles' Theorem we get:

**The Mean Value Theorem** Suppose $f(x)$ is continuous on $[a,b]$ and differentiable on $(a,b)$ then there is a $c \in (a,b)$ for which

$$f'(c) = \frac{f(b)-f(a)}{b-a}.$$




MVT says there is at least one c in $(a,b)$ with $f'(c) = \frac{f(b)-f(a)}{b-a}$ i.e. the two slopes are equal.

Read the proof of the MVT in the text, but also note and appreciate that the theorem is very intuitive and common-senseical.

Note that there can be several $c$ in $(a,b)$ with

$$f'(c) = \frac{f(b)-f(a)}{b-a}.$$

MVT says there is at least one such $c$.




**Example** (A way of thinking about the MVT.)

Suppose you drive 30 miles in 20 minutes. ($\frac{1}{3}$ hour). Did you break the speed limit? Your intuition say YES because your average velocity is $\frac{30 \text{ mi}}{\frac{1}{3} \text{ hour}} = 90$ mph. The mean value confirms this.

Say your position at time $t$ is $s(t)$


MVT says at some time $t=c$, $s'(c) = \underbrace{\frac{s(\frac{1}{3}) - s(0)}{\frac{1}{3} - 0}}_{\text{ave. vel.}} = \frac{30}{\frac{1}{3}} = 90$ mph

instantaneous velocity at time $c$

MVT simply says that at some instant $t=c$, your instantaneous velocity equals your average velocity

## Mathematical Consequences of MVT.

**Corollary 1** Suppose $f'(x) = 0$ on an interval $(a,b)$. Then $f(x) = C$ on $(a,b)$, where $C$ is a constant.




**Proof** Take $x \in (a,b)$. By MVT, there exists a $c \in [a, x]$ with $0 = f'(c) = \frac{f(a) - f(x)}{a - x}$

$\Rightarrow 0(a-x) = f(a) - f(x)$, i.e. $0 = f(a) - f(x)$. Then $f(x) = -f(a) = C$.

**Corollary 2** Suppose $f'(x) = g'(x)$ on $(a,b)$. Then $f(x) = g(x) + C$ for some constant $c$.




**Proof** If $f'(x) = g'(x)$ on $(a,b)$, then $f'(x) - g'(x) = 0$ on $(a,b)$. That is, $(f+g)'(x) = 0$ on $(a,b)$. By corollary 1, $(f+g)x = C$, i.e. $f(x) + g(x) = C$

Thus $f(x) = g(x) + (-C)$

Corollary says that if $f'(x) = g'(x)$, i.e. slopes of $f(x)$ and $g(x)$ agree then $f(x) = g(x) + C$