VCU

MATH 200

CALCULUS I

R. Hammack

Test 2

F

March 23, 2016

| Name: | | | | | |
|-------|--|--|--|--|--|
| - 10 | | | | | |
| | | | | | |

Score: _____

Directions. Answer the questions in the provided space. Unless noted otherwise, you must show and explain your work to receive full credit. Put your final answer in a box when appropriate.

This is a closed-book, closed-notes test. Calculators, computers, etc., are not used. Please put all phones away.

1. (20 points) Warmup: short answer.

(a) If
$$f(x) = \tan^{-1}(x) + \frac{1}{x}$$
, then $f'(x) =$

(b) If
$$f(x) = \tan(x) + e^x$$
, then $f'(x) =$

(c) If
$$f(x) = \sqrt[5]{x}^6$$
, then $f'(x) =$

(d) If
$$f(x) = \frac{1}{2}\sin(x) + e$$
, then $f'(x) =$

(e) If
$$f(x) = e^{\pi x}$$
, then $f'(x) =$

(f) If
$$f(x) = e^{\pi x}$$
, then $f'(0) =$

(g)
$$\lim_{h\to 0}\frac{e^{x+h}-e^x}{h}=$$

(h)
$$\frac{d}{dx} [\sin^{-1}(\pi x)] =$$

(i)
$$\frac{d}{dx} \left[\ln \left(\sin(x) \right) \right] =$$

(j)
$$\frac{d}{dx} \left[\frac{1}{x^2 + 3x} \right] =$$


2. (5 points) Find the equation of the tangent line to the graph of $y = \sin(x)$ at the point where $x = 2\pi$.

3. (5 points) Information about a function f(x) and its derivative is given in the table below.

| χ | 0 | 1 | 2 | 3 | 4 | 5 |
|---------------|----|----|----|---|----|----|
| f(x) | 0 | -3 | -2 | 3 | 10 | 25 |
| f(x) f'(x) | -1 | -7 | -5 | 5 | 20 | 30 |

Suppose $h(x) = f(x^2)$. Find h'(2). Show your work.

4. (5 points) A function f(x) is graphed below. Using the same coordinate axis, sketch the graph of the derivative f'(x).



5. (20 points) Find the following derivatives.

(a)
$$\frac{d}{dx} \left[\sec \left(e^{3x+1} \right) \right] =$$

(b)
$$\frac{d}{dx} \left[\ln \left(x^{10} - 4x^2 + 1 \right) \right] =$$

(c)
$$\frac{d}{dx} \left[\tan(x^5) + \tan^5(x) \right] =$$

(d)
$$\frac{d}{dx} \left[\frac{x^3 \ln(x)}{x^3 + 1} \right] =$$

6 (10 points) Find the derivative of $y = x^{\ln(x)}$

7 (10 points) Consider $f(x) = x^3 - 27x + 5$. Find all x for which the tangent to y = f(x) at the point (x, f(x)) is horizontal.

8. (15 points) An object moves on a straight line in such a way that its distance from its starting point at time t seconds is $s(t) = 4\sqrt{t}^5$ feet. What is its velocity is when its acceleration is 30 feet per second per second?

- **9.** (10 points) This question concerns the equation $x\sqrt[3]{y}^2 + y = 12$.
 - (a) Use implicit differentiation to find $\frac{dy}{dx}$.

(b) Use your answer from part (a) to find the slope of the tangent line to the graph of $x\sqrt[3]{y^2} + y = 12$ at the point (1,8).